

A.G. Kusraev and S.S. Kutateladze

**Some Aspects
of Boolean Valued Analysis**

УДК 517.98

Дата поступления 14 февраля 2015 г.

Kusraev A. G. and Kutateladze S. S.

SOME ASPECTS OF BOOLEAN VALUED ANALYSIS. — Новосибирск, 2015. — 19 с. — (Препринт / РАН. Сиб. отд-ние. Ин-т математики; No. 294).

This is an overview of some recent applications of Boolean valued analysis to operator theory and harmonic analysis. Under consideration are pseudoembedding operators, the noncommutative Wickstead problem, the Radon–Nikodým Theorem for JB -algebras, and the Bochner Theorem for lattice-valued positive definite mappings on locally compact groups.

KEYWORDS AND PHRASES: Boolean valued transfer principle, pseudoembedding operator, Wickstead problem, Radon–Nikodým theorem, Fourier transform, positive definite mapping, Bochner theorem.

Кусраев А. Г., Кутателадзе С. С.

НЕКОТОРЫЕ АСПЕКТЫ БУЛЕВОЗНАЧНОГО АНАЛИЗА

Обзор некоторых недавних приложений булевозначного анализа к теории операторов и гармоническому анализу. Рассмотрены псевдовложения, некоммутативная проблема Викстеда, теорема Радона — Никодима для JB -алгебр и теорема Бохнера для решеточнозначных положительно определенных отображений на локально компактных группах.

КЛЮЧЕВЫЕ СЛОВА И ФРАЗЫ: Булевозначный принцип переноса, псевдовложение, проблема Викстеда, теорема Радона — Никодима, преобразование Фурье, положительно определенное отображение, теорема Бохнера.

АДРЕСА АВТОРОВ:

Южный математический институт
ул. Маркуса, 22
362027 Владикавказ, Россия

Институт математики им. С. Л. Соболева СО РАН
пр. Академика Коптюга, 4
630090 Новосибирск, Россия

E-MAIL: kusraev@smath.ru; sskut@math.nsc.ru

- © Кусраев А. Г., Кутателадзе С. С., 2015
- © Южный математический институт, 2015
- © Институт математики им. С. Л. Соболева, 2015

Some Aspects of Boolean Valued Analysis

1. INTRODUCTION

We survey here some aspects of Boolean valued analysis that concern operator theory. The term *Boolean valued analysis* signifies the technique of studying properties of an arbitrary mathematical object by comparison between its representations in two different set-theoretic models whose construction utilizes principally distinct Boolean algebras. As these models, the classical Cantorian paradise in the shape of the von Neumann universe \mathbb{V} and a specially-trimmed Boolean valued universe $\mathbb{V}^{(\mathbb{B})}$ are usually taken. Comparison analysis is carried out by some interplay between \mathbb{V} and $\mathbb{V}^{(\mathbb{B})}$.

Boolean valued analysis not only is tied up with many topological and geometrical ideas but also provides a technology for expanding the content of the already available theorems. Each theorem, proven by the classical means, possesses some new unobvious content that relates to “variable sets.” A general scheme of the method is as follows; see [22, 23]. Assume that $\mathbf{X} \subset \mathbb{V}$ and $\mathcal{X} \subset \mathbb{V}^{(\mathbb{B})}$ are two classes of mathematical objects. Suppose that we are able to prove

The *Boolean Valued Representation*: Every $X \in \mathbf{X}$ embeds into a Boolean valued model, becoming an object $\mathcal{X} \in \mathcal{X}$ within $\mathbb{V}^{(\mathbb{B})}$.

The *Boolean Valued Transfer Principle* tells us then that every theorem about \mathcal{X} within Zermelo–Fraenkel set theory has its counterpart for the original object X interpreted as a Boolean valued object \mathcal{X} .

The *Boolean Valued Machinery* enables us to perform some translation of theorems from $\mathcal{X} \in \mathbb{V}^{(\mathbb{B})}$ to $X \in \mathbb{V}$ by using the appropriate general operations (ascending–descending) and the principles of Boolean valued analysis.

Everywhere below \mathcal{R} stands for the reals within $\mathbb{V}^{(\mathbb{B})}$. The Gordon Theorem states that the descent $\mathcal{R}\downarrow$ which is an algebraic system in \mathbb{V} is a universally complete vector lattice; see [22, 23]. Moreover, there exists an isomorphism χ of \mathbb{B} onto the Boolean algebra $\mathbb{P}(\mathcal{R}\downarrow)$ such that

$$\begin{aligned} \chi(b)x &= \chi(b)y \iff b \leq \llbracket x = y \rrbracket, \\ \chi(b)x \leq \chi(b)y &\iff b \leq \llbracket x \leq y \rrbracket \end{aligned} \tag{G}$$

for all $x, y \in \mathcal{R}\downarrow$ and $b \in \mathbb{B}$. The *restricted descent* $\mathcal{R}\downarrow$ of \mathcal{R} is the part of $\mathcal{R}\downarrow$ consisting of elements $x \in \mathcal{R}\downarrow$ with $|x| \leq C\mathbf{1}$ for some $C \in \mathbb{R}$, where $\mathbf{1}$ is an order unit in $\mathcal{R}\downarrow$.

By a *vector lattice* throughout the sequel we mean an Archimedean real vector lattice. We denote the Boolean algebras of all bands and all band projections in a vector lattice X respectively by $\mathbb{B}(X)$ and $\mathbb{P}(X)$ and we let $\mathcal{Z}(X)$ and $\text{Orth}(X)$ stand for the *ideal center* of X and the *f-algebra of orthomorphisms* on X , respectively. The universal completion X^u of a vector lattice X is always considered as a semiprime *f-algebra* whose multiplication is uniquely determined by fixing an order unit as a ring unit. The space of all order bounded linear operators from X to Y is denoted by $L^\sim(X, Y)$. The Riesz–Kantorovich Theorem tells us that if Y is a Dedekind complete vector lattice then so is $L^\sim(X, Y)$.

A linear operator T from X to Y is a *lattice homomorphism* whenever T preserves lattice operations; i.e., $T(x_1 \vee x_2) = Tx_1 \vee Tx_2$ (and so $T(x_1 \wedge x_2) = Tx_1 \wedge Tx_2$) for

all $x_1, x_2 \in X$. Vector lattices X and Y are said to be *lattice isomorphic* if there is a lattice isomorphism from X onto Y , i.e., T and T^{-1} are lattice homomorphisms. Let $\text{Hom}(X, Y)$ stand for the set of all lattice homomorphisms from X to Y . Recall also that the elements of the band $L_d^\sim(X, Y) := \text{Hom}(X, Y)^\perp$ are referred to as *diffuse operators*. An order bounded operator $T : X \rightarrow Y$ is said to be *pseudoembedding* if T belongs to the complementary band $L_a^\sim(X, Y) := \text{Hom}(X, Y)^{\perp\perp}$, the band generated by all lattice homomorphisms. Put $X^\sim := L^\sim(X, \mathbb{R})$ and $X_a^\sim := L_a^\sim(X, \mathbb{R})$.

We let $:=$ denote the assignment by definition, while \mathbb{N} , \mathbb{R} , and \mathbb{C} symbolize the naturals, the reals, and the complexes. Throughout the sequel \mathbb{B} is a complete Boolean algebra with top $\mathbb{1}$ and bottom $\mathbb{0}$. A *partition of unity* in \mathbb{B} is a family $(b_\xi)_{\xi \in \Xi}$ in \mathbb{B} such that $\bigvee_{\xi \in \Xi} b_\xi = \mathbb{1}$ and $b_\xi \wedge b_\eta = \mathbb{0}$ whenever $\xi \neq \eta$.

The reader can find the relevant information on the theory of order bounded operators in [6, 18, 27, 41, 45]; on the Boolean valued models of set theory, in [7, 39]; and on Boolean valued analysis, in [21, 22].

2. PSEUDOEMBEDDING OPERATORS

In this section we will give a description of the band generated by disjointness preserving operators in the vector lattice of order bounded operators. First we examine the scalar case.

2.1. *For an arbitrary vector lattice X there exist a unique cardinal γ and a disjoint family $(\varphi_\alpha)_{\alpha < \gamma}$ of nonzero lattice homomorphisms $\varphi_\alpha : X \rightarrow \mathbb{R}$ such that every $f \in X^\sim$ admits the unique representation*

$$f = f_d + o\text{-}\sum_{\alpha < \gamma} \lambda_\alpha \varphi_\alpha$$

where $f_d \in X_a^\sim$ and $(\lambda_\alpha)_{\alpha < \gamma} \subset \mathbb{R}$. The family $(\varphi_\alpha)_{\alpha < \gamma}$ is unique up to permutation and positive scalar multiplication.

◁ The Dedekind complete vector lattice X^\sim splits into the direct sum of the atomic band X_a^\sim and the diffuse band $X_d^\sim := (X_a^\sim)^\perp$; therefore, each functional $f \in E^\sim$ admits the unique representation $f = f_a + f_d$ with $f_a \in X_a^\sim$ and $f_d \in X_d^\sim$. Let γ be the cardinality of the set \mathcal{K} of one-dimensional bands in X_a^\sim (= atoms in $\mathbb{B}(X^\sim)$). Then there exists a family of lattice homomorphisms $(\varphi_\alpha : X \rightarrow \mathbb{R})_{\alpha < \gamma}$ such that $\mathcal{K} = \{\varphi_\alpha^\perp : \alpha < \gamma\}$. It remains to observe that the mapping sending a family of reals $(\lambda_\alpha)_{\alpha < \gamma}$ to the functional $x \mapsto o\text{-}\sum_{\alpha < \gamma} \lambda_\alpha \varphi_\alpha(x)$ implements a lattice isomorphism between X_a^\sim and some ideal in the vector lattice \mathbb{R}^γ .

If $(\psi_\alpha)_{\alpha < \gamma}$ is a disjoint family of nonzero real lattice homomorphisms on X then for all $\alpha, \beta < \gamma$ the functionals φ_α and ψ_β are either disjoint or proportional with a strictly positive coefficient, so that there exist a permutation $(\omega_\beta)_{\beta < \gamma}$ of $(\varphi_\alpha)_{\alpha < \gamma}$ and a unique family $(\mu_\beta)_{\beta < \gamma}$ in \mathbb{R}_+ such that $\psi_\beta = \mu_\beta \omega_\beta$ for all $\beta < \gamma$. ▷

2.2. Given two families $(S_\alpha)_{\alpha \in A}$ and $(T_\beta)_{\beta \in B}$ in $L^\sim(X, Y)$, say that $(S_\alpha)_{\alpha \in A}$ is a $\mathbb{P}(Y)$ -*permutation* of $(T_\beta)_{\beta \in B}$ whenever there exists a double family $(\pi_{\alpha, \beta})_{\alpha \in A, \beta \in B}$ in $\mathbb{P}(Y)$ such that $S_\alpha = \sum_{\beta \in B} \pi_{\alpha, \beta} T_\beta$ for all $\alpha \in A$, while $(\pi_{\alpha, \beta})_{\alpha \in A}$ and $(\pi_{\bar{\alpha}, \beta})_{\beta \in B}$ are partitions of unity in $\mathbb{B}(Y)$ for all $\bar{\alpha} \in A$ and $\bar{\beta} \in B$. It is easily seen that in case $Y = \mathbb{R}$ this amounts to saying that there is a bijection $\nu : A \rightarrow B$ with $S_\alpha = T_{\nu(\alpha)}$ for all $\alpha \in A$; i.e., $(S_\alpha)_{\alpha \in A}$ is a permutation of $(T_\beta)_{\beta \in B}$. We also say that $(S_\alpha)_{\alpha \in A}$ is an $\text{Orth}(Y)$ -multiple of $(T_\alpha)_{\alpha \in A}$ whenever there exists a family of orthomorphisms $(\pi_\alpha)_{\alpha \in A}$ in $\text{Orth}(Y)$ such that

$S_\alpha = \pi_\alpha T_\alpha$ for all $\alpha \in A$. In case $Y = \mathbb{R}$ we evidently get that S_α is a scalar multiple of T_α for all $\alpha \in A$.

Using the above notation, define the two mappings $\mathcal{S} : A \rightarrow X^{\wedge\downarrow}$ and $\mathcal{T} : B \rightarrow X^{\wedge\downarrow}$ by putting $\mathcal{S}(\alpha) := S_\alpha \uparrow$ ($\alpha \in A$) and $\mathcal{T}(\beta) := T_\beta \uparrow$ ($\beta \in B$). Recall that \uparrow signifies the modified ascent; see [23, 1.6.8].

2.3. Define the internal mappings $\tau, \sigma \in \mathbb{V}^{(\mathbb{B})}$ as $\sigma := \mathcal{S} \uparrow$ and $\tau := \mathcal{T} \uparrow$. Then $(\sigma(\alpha))_{\alpha \in A^\wedge}$ is a permutation of $(\tau(\beta))_{\beta \in B^\wedge}$ within $\mathbb{V}^{(\mathbb{B})}$ if and only if $(S_\alpha)_{\alpha \in A}$ is a $\mathbb{P}(Y)$ -permutation of $(T_\beta)_{\beta \in B}$.

\triangleleft Assume that $(\sigma(\alpha))_{\alpha \in A^\wedge}$ is a permutation of $(\tau(\beta))_{\beta \in B^\wedge}$ within $\mathbb{V}^{(\mathbb{B})}$. Then there is a bijection $\nu : B^\wedge \rightarrow A^\wedge$ such that $\sigma(\alpha) = \tau(\nu(\alpha))$ for all $(\alpha \in A^\wedge)$. By [23, 1.5.8] $\nu \downarrow$ is a function from A to $(B^\wedge) \downarrow = \text{mix}(\{\beta^\wedge : \beta \in B\})$. Thus, for each $\alpha \in A$ there exists a partition of unity $(b_{\alpha,\beta})_{\beta \in B}$ such that $\nu \downarrow(\alpha) = \text{mix}_{\beta \in B}(b_{\alpha,\beta} \beta^\wedge)$. Since $\nu \downarrow$ is injective, we have

$$\begin{aligned} \mathbb{1} &= \llbracket (\forall \alpha_1, \alpha_2 \in A^\wedge)(\nu(\alpha_1) = \nu(\alpha_2) \rightarrow \alpha_1 = \alpha_2) \rrbracket \\ &= \bigwedge_{\alpha_1, \alpha_2 \in A} \llbracket \nu(\alpha_1^\wedge) = \nu(\alpha_2^\wedge) \rightarrow \alpha_1^\wedge = \alpha_2^\wedge \rrbracket \\ &= \bigwedge_{\alpha_1, \alpha_2} \llbracket \nu \downarrow(\alpha_1) = \nu \downarrow(\alpha_2) \rrbracket \Rightarrow \llbracket \alpha_1^\wedge = \alpha_2^\wedge \rrbracket, \end{aligned}$$

and so $\llbracket \nu \downarrow(\alpha_1) = \nu \downarrow(\alpha_2) \rrbracket \leq \llbracket \alpha_1^\wedge = \alpha_2^\wedge \rrbracket$ for all $\alpha_1, \alpha_2 \in A$. Taking this inequality and the definition of $\nu \downarrow$ into account yields

$$\begin{aligned} b_{\alpha_1, \beta} \wedge b_{\alpha_2, \beta} &\leq \llbracket \nu \downarrow(\alpha_1) = \beta^\wedge \rrbracket \wedge \llbracket \nu \downarrow(\alpha_2) = \beta^\wedge \rrbracket \\ &\leq \llbracket \nu \downarrow(\alpha_1) = \nu \downarrow(\alpha_2) \rrbracket \leq \llbracket \alpha_1^\wedge = \alpha_2^\wedge \rrbracket, \end{aligned}$$

so that $\alpha_1 \neq \alpha_2$ implies $b_{\alpha_1, \beta} \wedge b_{\alpha_2, \beta} = 0$ because $x \neq y \iff \llbracket x^\wedge = y^\wedge \rrbracket = 0$ by [23, 1.4.5(2)]. At the same time, surjectivity of ν implies

$$\begin{aligned} \mathbb{1} &= \llbracket (\forall \beta \in B^\wedge)(\exists \alpha \in A^\wedge) \beta = \nu(\alpha) \rrbracket \\ &= \bigwedge_{\beta \in B} \bigvee_{\alpha \in A} \llbracket \beta^\wedge = \nu \downarrow(\alpha) \rrbracket = \bigwedge_{\beta \in B} \bigvee_{\alpha \in A} b_{\alpha, \beta}. \end{aligned}$$

It follows that $(b_{\alpha,\beta})_{\alpha \in A}$ is a partition of unity in \mathbb{B} for all $\beta \in B$. By the choice of ν it follows that $b_{\alpha,\beta} \leq \llbracket \sigma(\alpha^\wedge) = \tau(\beta^\wedge) \rrbracket$, because of the estimations

$$\begin{aligned} b_{\alpha,\beta} &\leq \llbracket \sigma(\alpha^\wedge) = \tau(\nu(\alpha^\wedge)) \rrbracket \wedge \llbracket \nu(\alpha^\wedge) = \beta^\wedge \rrbracket \\ &\leq \llbracket \sigma(\alpha^\wedge) = \tau(\beta^\wedge) \rrbracket = \llbracket \mathcal{S}(\alpha) = \mathcal{T}(\beta) \rrbracket. \end{aligned}$$

Put $\pi_{\alpha,\beta} := \chi(b_{\alpha,\beta})$ and observe that $b_{\alpha,\beta} \leq \llbracket \mathcal{S}(\alpha)x^\wedge = \mathcal{T}(\beta)x^\wedge \rrbracket \leq \llbracket S_\alpha x = T_\beta x \rrbracket$ for all $\alpha \in A, \beta \in B$, and $x \in X$. Using (\mathbb{G}) , we obtain $\pi_{\alpha,\beta} S_\alpha = \pi_{\alpha,\beta} T_\beta$ and so $S_\alpha = \sum_{\beta \in B} \pi_{\alpha,\beta} T_\beta$ for all $\alpha \in A$. Clearly, $(\pi_{\alpha,\beta})$ is the family as required in 2.2. The sufficiency is shown by the same reasoning in the reverse direction. \triangleright

2.4. A nonempty set \mathcal{D} of positive operators from X to Y is called *strongly generating* if \mathcal{D} is a disjoint set and $S(X)^{\perp\perp} = Y$ for all $S \in \mathcal{D}$. If, in addition, $\mathcal{D}^{\perp\perp} = B$, then we say also that \mathcal{D} *strongly generates* the band $B \subset L^\sim(X, Y)$ or B is strongly generated by \mathcal{D} . In case $Y = \mathbb{R}$, the strongly generating sets in $X^\sim = L^\sim(X, \mathbb{R})$ are precisely disjoint sets of nonzero positive functionals.

Given a cardinal γ and a universally complete vector lattice Y , say that a vector lattice X is (γ, Y) -*homogeneous* if the band $L_a^\sim(X, Y)$ is strongly generated by a set of lattice

homomorphisms of cardinality γ and for every nonzero projection $\pi \in \mathbb{P}(Y)$ and every strongly generating set \mathcal{D} in $L_a^\sim(X, \pi Y)$ we have $\text{card}(\mathcal{D}) \geq \gamma$. We say also that X is (γ, π) -homogeneous if $\pi \in \mathbb{P}(Y)$ and X is $(\gamma, \pi Y)$ -homogeneous. Evidently, the (γ, \mathbb{R}) -homogeneity of a vector lattice X amounts just to saying that the band X_a^\sim is generated in X^\sim by a cardinality γ disjoint set of nonzero lattice homomorphisms or, equivalently, the cardinality of the set of atoms in $\mathbb{B}(X^\sim)$ equals γ .

Take $\mathcal{D} \subset L^\sim(X, \mathcal{R}\downarrow)$ and $\Delta \in \mathbb{V}^{(\mathbb{B})}$ with $\llbracket \Delta \subset (X^\wedge)^\sim \rrbracket = \mathbf{1}$. Put $\mathcal{D}_\uparrow := \{T\uparrow : T \in \mathcal{D}\}$ and $\Delta^\downarrow := \{\tau\downarrow : \tau \in \Delta\}$. Let $\text{mix}(\mathcal{D})$ stand for the set of all $T \in L^\sim(X, \mathcal{R}\downarrow)$ representable as $Tx = \sigma\text{-}\sum_{\xi \in \Xi} \pi_\xi T_\xi x$ ($x \in X$) with $(\pi_\xi)_{\xi \in \Xi}$ a partition of unity in $\mathbb{P}(\mathcal{R}\downarrow)$ and $(T_\xi)_{\xi \in \Xi}$ a family in \mathcal{D} .

2.5. *Let $\Delta \subset (X^\wedge)^\sim$ be a disjoint set of nonzero positive functionals which has cardinality γ^\wedge within $\mathbb{V}^{(\mathbb{B})}$. Then there exists a cardinality γ strongly generating set of positive operators \mathcal{D} from X to $\mathcal{R}\downarrow$ such that $\Delta = \mathcal{D}_\uparrow$ and $\Delta^\downarrow = \text{mix}(\mathcal{D})$.*

\triangleleft If Δ obeys the conditions then there is $\phi \in \mathbb{V}^{(\mathbb{B})}$ such that $\llbracket \phi : \gamma^\wedge \rightarrow \Delta \text{ is a bijection} \rrbracket = \mathbf{1}$. Note that $\varphi\downarrow$ sends γ into $\Delta\downarrow \subset (X^\wedge)^\sim\downarrow$ by [23, 1.5.8]. By [23, Theorem 3.3.3], we can define the mapping $\alpha \mapsto \Phi(\alpha)$ from γ to $L^\sim(X, \mathcal{R}\downarrow)$ by letting $\Phi(\alpha) := (\phi\downarrow(\alpha))\downarrow$. Put $\mathcal{D} := \{\Phi(\alpha) : \alpha \in \gamma\}$ and note that $\mathcal{D} \subset \Delta^\downarrow$. Using [23, 1.6.6] and the surjectivity of ϕ we have $\Delta\downarrow = \varphi(\gamma^\wedge)\downarrow = \text{mix}\{\phi\downarrow(\alpha) : \alpha \in \gamma\}$ and combining this with [23, 3.3.7] we get $\Delta = \mathcal{D}_\uparrow$ and $\Delta^\downarrow = \text{mix}(\mathcal{D})$.

The injectivity of ϕ implies that

$$\llbracket (\forall \alpha, \beta \in \gamma^\wedge)(\alpha \neq \beta \rightarrow \phi(\alpha) \neq \phi(\beta)) \rrbracket = \mathbf{1}.$$

Replacing the universal quantifier by the supremum over $\alpha, \beta \in \gamma^\wedge$, from [23, 1.4.5 (1) and 1.4.5 (2)] we deduce that

$$\mathbf{1} = \bigwedge_{\alpha, \beta \in \gamma} \llbracket \alpha^\wedge \neq \beta^\wedge \rrbracket \Rightarrow \llbracket \varphi(\alpha^\wedge) \neq \phi(\beta)^\wedge \rrbracket = \bigwedge_{\substack{\alpha, \beta \in \gamma \\ \alpha \neq \beta}} \llbracket \Phi(\alpha) \neq \Phi(\beta) \rrbracket,$$

and so $\alpha \neq \beta$ implies $\Phi(\alpha) \neq \Phi(\beta)$ for all $\alpha, \beta \in \gamma$. Thus Φ is injective and the cardinality of \mathcal{D} is γ . The fact that \mathcal{D} is strongly generating follows from [23, 3.3.5 (5) and 3.8.4]. \triangleright

2.6. *If \mathcal{D} is a cardinality γ strongly generating set of positive operators from X to $\mathcal{R}\downarrow$ then $\Delta = \mathcal{D}_\uparrow \subset (X^\wedge)^\sim$ is a disjoint set of nonzero positive functionals which has cardinality $|\gamma^\wedge|$ within $\mathbb{V}^{(\mathbb{B})}$.*

\triangleleft Assume that $\mathcal{D} \subset L(X, \mathcal{R}\downarrow)$ is a strongly generating set of cardinality γ . Then there is a bijection $f : \gamma \rightarrow \mathcal{D}_\uparrow$. Moreover, $\alpha \neq \beta$ implies $\llbracket f(\alpha) \perp f(\beta) \rrbracket = \mathbf{1}$ by [23, 3.3.5 (5)] and $\llbracket f(\alpha) \neq 0 \rrbracket = \mathbf{1}$ by [23, 3.8.4]. Interpreting in $\mathbb{V}^{(\mathbb{B})}$ the ZFC-theorem

$$(\forall f, g \in X^\sim)(f \neq 0 \wedge g \neq 0 \wedge f \perp g \rightarrow f \neq g)$$

yields $\llbracket f(\alpha) \neq f(\beta) \rrbracket = \mathbf{1}$ for all $\alpha, \beta \in \gamma$, $\alpha \neq \beta$. It follows that $\phi := f\uparrow$ is a bijection from γ^\wedge onto $\Delta = (\mathcal{D}_\uparrow)\uparrow$, so that the cardinality of Δ is $|\gamma^\wedge|$. The proof is completed by the arguments similar to those in 2.5. \triangleright

2.7. *A vector lattice X is $(\gamma, \mathcal{R}\downarrow)$ -homogeneous for some cardinal γ if and only if $\llbracket \gamma^\wedge \text{ is a cardinal and } X^\wedge \text{ is } (\gamma^\wedge, \mathcal{R})\text{-homogeneous} \rrbracket = \mathbf{1}$.*

\triangleleft *Sufficiency:* Assume that γ^\wedge is a cardinal and X^\wedge is $(\gamma^\wedge, \mathcal{R})$ -homogeneous within $\mathbb{V}^{(\mathbb{B})}$. The latter means that $(X^\wedge)_a^\sim$ is generated by a disjoint set of nonzero lattice homomorphisms $\Delta \subset (X^\wedge)^\sim$ which has cardinality γ^\wedge within $\mathbb{V}^{(\mathbb{B})}$. By 2.5 there exists a strongly generating set \mathcal{D} in $L_a^\sim(X, \mathcal{R}\downarrow)$ of cardinality γ^\wedge such that $\Delta = \mathcal{D}_\uparrow$. Take a nonzero $\pi \in \mathbb{P}(\mathcal{R}\downarrow)$

and put $b := \chi^{-1}(\pi)$. Recall that we can identify $L^\sim(X, \pi(\mathcal{R}\downarrow))$ and $L^\sim(X, (b \wedge \mathcal{R})\downarrow)$. If \mathcal{D}' is a strongly generating set in $L_a^\sim(X, \pi(\mathcal{R}\downarrow))$ of cardinality β then \mathcal{D}'_\uparrow strongly generates $(X^\wedge)_a^\sim$ and has cardinality $|\beta^\wedge|$ within the relative universe $\mathbb{V}^{([0, b])}$. By [23, 1.3.7] $\gamma^\wedge = |\beta^\wedge| \leq \beta^\wedge$ and so $\gamma \leq \beta$.

Necessity: Assume now that X is $(\gamma, \mathcal{R}\downarrow)$ -homogeneous and the set \mathcal{D} of lattice homomorphisms of cardinality γ strongly generates the band $L_a^\sim(X, \mathcal{R}\downarrow)$. Then $\Delta = \mathcal{D}_\uparrow$ generates the band $(X^\wedge)_a^\sim$ and the cardinalities of Δ and γ^\wedge coincide; i.e., $|\Delta| = |\gamma^\wedge|$. By [23, 1.9.11] the cardinal $|\gamma^\wedge|$ has the representation $|\gamma^\wedge| = \text{mix}_{\alpha \leq \gamma} b_\alpha \alpha^\wedge$, where $(b_\alpha)_{\alpha \leq \gamma}$ is a partition of unity in \mathbb{B} . It follows that $b_\alpha \leq \llbracket \Delta \text{ is a generating set in } (X^\wedge)_a^\sim \text{ of cardinality } \alpha^\wedge \rrbracket = \mathbb{1}$. If $b_\alpha \neq 0$ then $b_\alpha \wedge \Delta$ is a generating set in $(X^\wedge)_a^\sim$ of cardinality $|\gamma^\wedge| = \alpha^\wedge \leq \gamma^\wedge$ in the relative universe $\mathbb{V}^{[0, b_\alpha]}$. Put $\pi_\alpha = \chi(b_\alpha)$ and $\pi_\alpha \circ \mathcal{D} := \{\pi_\alpha \circ T : T \in \mathcal{D}\}$. Clearly, $b_\alpha \wedge \Delta = (\pi_\alpha \mathcal{D})_\uparrow$ and so $\pi_\alpha \circ \mathcal{D}$ strongly generates the band $L_a^\sim(X, \mathcal{R}\downarrow)$. By hypothesis \mathcal{D} is $(\gamma, \mathcal{R}\downarrow)$ -homogeneous. Consequently, $\alpha \geq \gamma$, so that $\alpha = \gamma$, since $\alpha \leq \gamma$ if and only if $\alpha^\wedge \leq \gamma^\wedge$. Thus, $|\gamma^\wedge| = \gamma^\wedge$ whenever $b_\alpha \neq 0$ and γ^\wedge is a cardinal within $\mathbb{V}^{(\mathbb{B})}$. \triangleright

2.8. *Let X be a (γ, Y) -homogeneous vector lattice for some universally complete vector lattice Y and a nonzero cardinal γ . Then there exists a strongly generating family of lattice homomorphisms $(\Phi_\alpha)_{\alpha < \gamma}$ from X to Y such that each operator $T \in L_a^\sim(X, Y)$ admits the unique representation $T = o\text{-}\sum_{\alpha < \gamma} \sigma_\alpha \circ \Phi_{\gamma, \alpha}$, where $(\sigma_\alpha)_{\alpha < \gamma}$ is a family of orthomorphisms in $\text{Orth}(Y)$.*

\triangleleft This is immediate from the definitions in 2.4. \triangleright

2.9. Theorem. *Let X and Y be vector lattices with Y universally complete. Then there are a nonempty set of cardinals Γ and a partition of unity $(Y_\gamma)_{\gamma \in \Gamma}$ in $\mathbb{B}(Y)$ such that X is (γ, Y_γ) -homogeneous for all $\gamma \in \Gamma$.*

\triangleleft We may assume without loss of generality that $Y = \mathcal{R}\downarrow$. The transfer principle tells us that according to 2.1 there exists a cardinal \varkappa within $\mathbb{V}^{(\mathbb{B})}$ such that $(X^\wedge)_a^\sim$ is generated by a cardinality \varkappa disjoint set \mathcal{H} of nonzero \mathbb{R}^\wedge -linear lattice homomorphisms or, equivalently, $\llbracket X^\wedge \text{ is } (\varkappa, \mathcal{R})\text{-homogeneous} \rrbracket = \mathbb{1}$. By [23, 1.9.11] there is a nonempty set of cardinals Γ and a partition of unity $(b_\gamma)_{\gamma \in \Gamma}$ in \mathbb{B} such that $\varkappa = \text{mix}_{\gamma \in \Gamma} b_\gamma \gamma^\wedge$. It follows that $b_\gamma \leq \llbracket X^\wedge \text{ is } (\gamma^\wedge, \mathcal{R})\text{-homogeneous} \rrbracket$ for all $\gamma \in \Gamma$. Passing to the relative subalgebra $\mathbb{B}_\gamma := [0, b_\gamma]$ and considering [23, 1.3.7] we conclude that $\mathbb{V}^{(\mathbb{B}_\gamma)} \models \text{“}X^\wedge \text{ is } (\gamma^\wedge, b_\gamma \wedge \mathcal{R})\text{-homogeneous”}$, so that X is $(\gamma, (b_\gamma \wedge \mathcal{R})\downarrow)$ -homogeneous by 2.7. In view of [23, 2.3.6] $(b_\gamma \wedge \mathcal{R})\downarrow$ is lattice isomorphic to Y_γ , and so the desired result follows. \triangleright

2.10. Theorem. *Let X and Y be vector lattices with Y universally complete. Then there are a nonempty set of cardinals Γ and a partition of unity $(Y_\gamma)_{\gamma \in \Gamma}$ in $\mathbb{B}(Y)$ such that to each cardinal $\gamma \in \Gamma$ there is a disjoint family of lattice homomorphisms $(\Phi_{\gamma, \alpha})_{\alpha < \gamma}$ from X to Y_γ satisfying*

(1) $\Phi_{\gamma, \alpha}(X)^{\perp\perp} = Y_\gamma \neq \{0\}$ for all $\gamma \in \Gamma$ and $\alpha < \gamma$.

(2) X is (γ, Y_γ) -homogeneous for all $\gamma \in \Gamma$.

(3) For each order dense sublattice $Y_0 \subset Y$ each $T \in L^\sim(X, Y_0)$ admits the unique representation

$$T = T_d + o\text{-}\sum_{\gamma \in \Gamma} o\text{-}\sum_{\alpha < \gamma} \sigma_{\gamma, \alpha} \circ \Phi_{\gamma, \alpha},$$

with $T_d \in L_a^\sim(X, Y)$ and $\sigma_{\gamma, \alpha} \in \text{Orth}(\Phi_{\gamma, \alpha}, Y_0)$.

For every $\gamma \in \Gamma$ the family $(\Phi_{\gamma,\alpha})_{\alpha < \gamma}$ is unique up to $\mathbb{P}(Y)$ -permutation and $\text{Orth}(Y_\gamma)_+$ -multiplication.

◁ The existence of $(Y_\gamma)_{\gamma \in \Gamma}$ and $(\Phi_{\gamma,\alpha})_{\gamma \in \Gamma, \alpha < \gamma}$ with the required properties is immediate from 2.8 and 2.9. The uniqueness follows from 2.1 and 2.3. ▷

2.11. Theorem 2.10, the main result of Section 2, was proved by Tabuev in [35, Theorem 2.2] with standard tools. The pseudoembedding operators are closely connected with the so-called order narrow operators. A linear operator $T : X \rightarrow Y$ is *order narrow* if for every $x \in X_+$ there exists a net (x_α) in X such that $|x_\alpha| = x$ for all α and (Tx_α) is order convergent to zero in Y ; see [28, Definition 3.1]. The main result by Maslyuchenko, Mykhaylyuk, and Popov in [28, Theorem 11.7(ii)] states that if X and Y are Dedekind complete vector lattices with X atomless and Y an order ideal of some order continuous Banach lattice then an order bounded order continuous operator is order narrow if and only if it is pseudoembedding.

2.12. The term *pseudoembedding operator* stems from a result by Rosenthal [31] which asserts that a nonzero bounded linear operator in L^1 is a pseudoembedding if and only if it is a near isometric embedding when restricted to a suitable $L^1(A)$ -subspace. Systematic study of narrow operators was started by Plichko and Popov in [29]. Concerning a detailed presentation of the theory of narrow operators, we refer to the recent book by Popov and Randrianantoanina [30] and the references therein.

3. THE NONCOMMUTATIVE WICKSTEAD PROBLEM

When are we so happy in a vector lattice that all band preserving linear operators turn out to be order bounded?

This question was raised by Wickstead in [42]. We refer to [22, 23] and [13] for a detailed presentation of the Wickstead problem. In this section we consider a noncommutative version of the problem. The relevant information on the theory of Baer $*$ -algebras and AW^* -algebras can be found in Berberian [9], Chilin [10], and Kusraev [18].

3.1. A *Baer $*$ -algebra* is a complex involutive algebra A such that, for each nonempty $M \subset A$, there is a projection, i.e., a hermitian idempotent p , satisfying $M^\perp = pA$ where $M^\perp := \{y \in A : (\forall x \in M) xy = 0\}$ is the right annihilator of M . Clearly, this amounts to saying that each left annihilator has the form ${}^\perp M = Aq$ for an appropriate projection q . To each left annihilator L in a Baer $*$ -algebra there is a unique projection $q_L \in A$ such that $x = xq_L$ for all $x \in L$ and $q_L y = 0$ whenever $y \in L^\perp$. The mapping $L \mapsto q_L$ is an isomorphism between the poset of left annihilators and the poset of all projections. Thus, the poset $\mathbb{P}(A)$ of all projections in a Baer $*$ -algebra is an order complete lattice. (Clearly, the formula $q \leq p \iff q = qp = pq$, sometimes pronounced as “ p contains q ,” specifies some order on the set of projections $\mathbb{P}(A)$.)

An element z in A is *central* provided that z commutes with every member of A ; i.e., $(\forall x \in A) xz = zx$. The *center* of a Baer $*$ -algebra A is the set $\mathcal{Z}(A)$ comprising central elements. Clearly, $\mathcal{Z}(A)$ is a commutative Baer $*$ -subalgebra of A , with $\lambda \mathbb{1} \in \mathcal{Z}(A)$ for all $\lambda \in \mathbb{C}$. A *central projection* of A is a projection belonging to $\mathcal{Z}(A)$. Put $\mathbb{P}_c(A) := \mathbb{P}(A) \cap \mathcal{Z}(A)$.

3.2. A *derivation* on a Baer $*$ -algebra A is a linear operator $d : A \rightarrow A$ satisfying $d(xy) = d(x)y + xd(y)$ for all $x, y \in A$. A derivation d is *inner* provided that $d(x) = ax - xa$

($x \in A$) for some $a \in A$. Clearly, an inner derivation vanishes on $\mathcal{Z}(A)$ and is $\mathcal{Z}(A)$ -linear; i.e., $d(ex) = ed(x)$ for all $x \in A$ and $e \in \mathcal{Z}(A)$.

Consider a derivation $d : A \rightarrow A$ on a Baer $*$ -algebra A . If $p \in A$ is a central projection then $d(p) = d(p^2) = 2pd(p)$. Multiplying this identity by p we have $pd(p) = 2pd(p)$ so that $d(p) = pd(p) = 0$. Consequently, every derivation vanishes on the linear span of $\mathbb{P}_c(A)$, the set of all central projections. In particular, $d(ex) = ed(x)$ whenever $x \in A$ and e is a linear combination of central projections. Even if the linear span of central projections is dense in a sense in $\mathcal{Z}(A)$, the derivation d may fail to be $\mathcal{Z}(A)$ -linear.

This brings up the natural question: *Under what conditions is every derivation Z -linear on a Baer $*$ -algebra A provided that Z is a Baer $*$ -subalgebra of $\mathcal{Z}(A)$?*

3.3. An AW^* -algebra is a C^* -algebra with unity $\mathbb{1}$ which is also a Baer $*$ -algebra. More explicitly, an AW^* -algebra is a C^* -algebra whose every right annihilator has the form pA , with p a projection. Clearly, $\mathcal{Z}(A)$ is a commutative AW^* -subalgebra of A . If $\mathcal{Z}(A) = \{\lambda\mathbb{1} : \lambda \in \mathbb{C}\}$ then the AW^* -algebra A is an AW^* -factor.

3.4. A C^* -algebra A is an AW^* -algebra if and only if the two conditions hold:

(1) Each orthogonal family in $\mathbb{P}(A)$ has a supremum.

(2) Each maximal commutative $*$ -subalgebra of $A_0 \subset A$ is a Dedekind complete f -algebra (or, equivalently, coincides with the least norm closed $*$ -subalgebra containing all projections of A_0).

3.5. Given an AW^* -algebra A , define the two sets $C(A)$ and $S(A)$ of measurable and locally measurable operators, respectively. Both are Baer $*$ -algebras; cp. Chilin [10]. Suppose that Λ is an AW^* -subalgebra in $\mathcal{Z}(A)$, and Φ is a Λ -valued trace on A_+ . Then we can define another Baer $*$ -algebra, $L(A, \Phi)$, of Φ -measurable operators. The center $\mathcal{Z}(A)$ is a vector lattice with a strong unit, while the centers of $C(A)$, $S(A)$, and $L(A, \Phi)$ coincide with the universal completion of $\mathcal{Z}(A)$. If d is a derivation on $C(A)$, $S(A)$, or $L(A, \Phi)$ then $d(px) = pd(x)$ ($p \in \mathbb{P}_c(A)$) so that d can be considered as band preserving in a sense (cp. [23, 4.1.1 and 4.10.4]). The natural question arises immediately about these algebras:

3.6. $WP(C)$: *When are all derivations on $C(A)$, $S(A)$, or $L(A, \Phi)$ inner?* This question may be regarded as the *noncommutative Wickstead problem*.

3.7. The classification of AW^* -algebras into types is determined from their lattices of projections $\mathbb{P}(A)$; see Sakai [32]. We recall only the definition of type I AW^* -algebra. A projection $\pi \in A$ is *abelian* if $\pi A \pi$ is a commutative algebra. An algebra A has *type I* provided that each nonzero projection in A contains a nonzero abelian projection.

A C^* -algebra A is \mathbb{B} -embeddable provided that there are a type I AW^* -algebra N and a $*$ -monomorphism $\iota : A \rightarrow N$ such that $\mathbb{B} = \mathbb{P}_c(N)$ and $\iota(A) = \iota(A)''$, where $\iota(A)''$ is the bicommutant of $\iota(A)$ in N . Note that in this event A is an AW^* -algebra and \mathbb{B} is a complete subalgebra of $\mathbb{P}_c(A)$.

3.8. Theorem. *Let A be a type I AW^* -algebra, let Λ be an AW^* -subalgebra of $\mathcal{Z}(A)$, and let Φ be a Λ -valued faithful normal semifinite trace on A . If the complete Boolean algebra $\mathbb{B} := \mathbb{P}(\Lambda)$ is σ -distributive and A is \mathbb{B} -embeddable, then every derivation on $L(A, \Phi)$ is inner.*

\triangleleft We briefly sketch the proof. Let $\mathcal{A} \in \mathbb{V}^{(\mathbb{B})}$ be the Boolean valued representation of A . Then \mathcal{A} is a von Neumann algebra within $\mathbb{V}^{(\mathbb{B})}$. Since the Boolean valued interpretation preserves classification into types, \mathcal{A} is of type I. Let φ stand for the Boolean valued

representation of Φ . Then φ is a faithful normal semifinite \mathcal{C} -valued trace on \mathcal{A} and the descent of $L(\mathcal{A}, \varphi)$ is $*$ - Λ -isomorphic to $L(A, \Phi)$; cp. Korol' and Chilin [17]. Suppose that d is a derivation on $L(A, \Phi)$ and δ is the Boolean valued representation of d . Then δ is a \mathcal{C} -valued \mathbb{C}^\wedge -linear derivation on $L(\mathcal{A}, \varphi)$. Since \mathbb{B} is σ -distributive, $\mathcal{C} = \mathbb{C}^\wedge$ within $\mathbb{V}^{(\mathbb{B})}$ and δ is \mathcal{C} -linear. But it is well known that every derivation on a type I von Neumann algebra is inner; cp. [4]. Therefore, d is also inner. \triangleright

3.8. Theorem 3.8 is taken from Gutman, Kusraev, and Kutateladze [13, Theorem 4.3.6]. This fact is an interesting ingredient of the theory of noncommutative integration which stems from Segal [33]. Considerable attention is given to derivations on various algebras of measurable operators associated with an AW^* -algebra and a central-valued trace. We mention only the article [4] by Albeverio, Ajupov, and Kudaybergenov and the article [8] by Ber, de Pagter, and Sukochev.

4. THE RADON–NIKODÝM THEOREM FOR JB -ALGEBRAS

In this section we sketch some further applications of the Boolean value approach to nonassociative Radon–Nikodým type theorems.

4.1. Let A be a vector space over some field \mathbb{F} . Say that A is a *Jordan algebra*, if there is given a (generally) nonassociative binary operation $A \times A \ni (x, y) \mapsto xy \in A$ on A , called *multiplication* and satisfying the following for all $x, y, z \in A$ and $\alpha \in \mathbb{F}$:

- (1) $xy = yx$;
- (2) $(x + y)z = xz + yz$;
- (3) $\alpha(xy) = (\alpha x)y$;
- (4) $(x^2y)x = x^2(yx)$.

An element e of a Jordan algebra A is a *unit element* or a *unit* of A , if $e \neq 0$ and $ea = a$ for all $a \in A$.

4.2. Recall that a *JB-algebra* is simultaneously a real Banach space A and a unital Jordan algebra with unit $\mathbb{1}$ such that

- (1) $\|xy\| \leq \|x\| \cdot \|y\| \quad (x, y \in A)$,
- (2) $\|x^2\| = \|x\|^2 \quad (x \in A)$,
- (3) $\|x^2\| \leq \|x^2 + y^2\| \quad (x, y \in A)$.

The set $A_+ := \{x^2 : x \in A\}$, presenting a proper convex cone, determines the structure of an ordered vector space on A so that the unity $\mathbb{1}$ of the algebra A serves as a strong order unit, and the order interval $[-\mathbb{1}, \mathbb{1}] := \{x \in A : -\mathbb{1} \leq x \leq \mathbb{1}\}$ serves as the unit ball. Moreover, the inequalities $-\mathbb{1} \leq x \leq \mathbb{1}$ and $0 \leq x^2 \leq \mathbb{1}$ are equivalent.

The intersection of all maximal associative subalgebras of A is called the *center* of A and denoted by $\mathcal{Z}(A)$. The element a belongs to $\mathcal{Z}(A)$ if and only if $(ax)y = a(xy)$ for all $x, y \in A$. If $\mathcal{Z}(A) = \mathbb{R} \cdot \mathbb{1}$, then A is said to be a *JB-factor*. The center $Z(A)$ is an associative *JB-algebra*, and such an algebra is isometrically isomorphic to the real Banach algebra $C(Q)$ of continuous functions on some compact space Q .

4.3. The idempotents of a *JB-algebra* are also called *projections*. The set of all projections $\mathbb{P}(A)$ forms a complete lattice with the order defined as $\pi \leq \rho \iff \pi \circ \rho = \pi$. The sublattice of *central projections* $\mathbb{P}_c(A) := \mathbb{P}(A) \cap \mathcal{Z}(A)$ is a Boolean algebra. Assume that \mathbb{B} is a subalgebra of the Boolean algebra $\mathbb{P}_c(A)$ or, equivalently, $\mathbb{B}(\mathbb{R})$ is a subalgebra

of the center $\mathcal{Z}(A)$ of A . Then we say that A is a \mathbb{B} -*JB-algebra* if, for every partition of unity $(e_\xi)_{\xi \in \Xi}$ in \mathbb{B} and every family $(x_\xi)_{\xi \in \Xi}$ in A , there exists a unique \mathbb{B} -mixing $x := \text{mix}_{\xi \in \Xi}(e_\xi x_\xi)$, i.e., a unique element $x \in A$ such that $e_\xi x_\xi = e_\xi x$ for all $\xi \in \Xi$. If $\mathbb{B}(\mathbb{R}) = \mathcal{Z}(A)$, then a \mathbb{B} -*JB-algebra* is also referred to as *centrally extended JB-algebra*.

The unit ball of a \mathbb{B} -*JB-algebra* is closed under \mathbb{B} -mixings. Consequently, each \mathbb{B} -*JB-algebra* is a \mathbb{B} -cyclic Banach space.

4.4. Theorem. *The restricted descent of a JB-algebra in the model $\mathbb{V}^{(\mathbb{B})}$ is a \mathbb{B} -JB-algebra. Conversely, for every \mathbb{B} -JB-algebra A there exists a unique (up to isomorphism) JB-algebra \mathcal{A} within $\mathbb{V}^{\mathbb{B}}$ whose restricted descent is isometrically \mathcal{B} -isomorphic to A . Moreover, $\llbracket \mathcal{A} \text{ is a JB-factor} \rrbracket = \mathbf{1}$ if and only if $\mathbb{B}(\mathbb{R}) = \mathcal{Z}(A)$.*

◁ See [22, Theorem 12.7.6] and [19, Theorem 3.1]. ▷

4.5. Now we give two applications of the above Boolean valued representation result to \mathbb{B} -*JB-algebras*. Theorems 4.7 and 4.11 below appear by transfer of the corresponding facts from the theory of *JB-algebras*.

Let A be a \mathbb{B} -*JB-algebra* and $\Lambda := \mathbb{B}(\mathbb{R})$. An operator $\Phi \in A^\sharp$ is called a Λ -*valued state* if $\Phi \geq 0$ and $\Phi(\mathbf{1}) = \mathbf{1}$. A state Φ is said to be *normal* if, for every increasing net (x_α) in A with the least upper bound $x := \sup x_\alpha$, we have $\Phi(x) = o\text{-}\lim \Phi(x_\alpha)$. If \mathcal{A} is the Boolean valued representation of the algebra A , then the ascent $\varphi := \Phi \uparrow$ is a bounded linear functional on \mathcal{A} by [23, Theorem 5.8.12]. Moreover, φ is positive and order continuous; i.e., φ is a normal state on \mathcal{A} . The converse is also true: if $\llbracket \varphi \text{ is a normal state on } \mathcal{A} \rrbracket = \mathbf{1}$, then the restriction of the operator $\varphi \downarrow$ to A is a Λ -valued normal state. Now we will characterize \mathbb{B} -*JB-algebras* that are \mathbb{B} -dual spaces. Toward this end, it suffices to give Boolean valued interpretation for the following result.

4.6. Theorem. *A JB-algebra is a dual Banach space if and only if it is monotone complete and has a separating set of normal states.*

◁ See [34, Theorem 2.3]. ▷

4.7. Theorem. *Let \mathbb{B} be a complete Boolean algebra and let Λ be a Dedekind complete unital AM-space with $\mathbb{B} \simeq \mathbb{P}(\Lambda)$. A \mathbb{B} -JB-algebra A is a \mathbb{B} -dual space if and only if A is monotone complete and admits a separating set of Λ -valued normal states. If one of these equivalent conditions holds, then the part of A^\sharp consisting of order continuous operators serves as a \mathbb{B} -predual space of A .*

◁ See [22, Theorem 12.8.5] and [19, Theorem 4.2]. ▷

4.8. An algebra A satisfying one of the equivalent conditions 4.7 is called a \mathbb{B} -*JBW-algebra*. If, moreover, \mathbb{B} coincides with the set of all central projections, then A is said to be a \mathbb{B} -*JBW-factor*. It follows from Theorems 4.4 and 4.7 that A is a \mathbb{B} -*JBW-algebra* (\mathbb{B} -*JBW-factor*) if and only if its Boolean valued representation $\mathcal{A} \in V^{(\mathbb{B})}$ is a *JBW-algebra* (*JBW-factor*).

A mapping $\Phi : A_+ \rightarrow \Lambda \cup \{+\infty\}$ is a (Λ -valued) *weight* if the conditions are satisfied (under the assumptions that $\lambda + (+\infty) := +\infty + \lambda := +\infty$, $\lambda \cdot (+\infty) := \lambda$ for all $\lambda \in \Lambda$, while $0 \cdot (+\infty) := 0$ and $+\infty + (+\infty) := +\infty$):

(1) $\Phi(x + y) = \phi(x) + \Phi(y)$ for all $x, y \in A_+$.

(2) $\Phi(\lambda x) = \lambda \Phi(x)$ for all $x \in A_+$ and $\lambda \in \Lambda_+$.

A weight Φ is said to be a *trace* if the additional condition is satisfied

(3) $\Phi(x) = \Phi(U_s x)$ for all $x \in A_+$ and $s \in A$ with $s^2 = \mathbf{1}$.

Here U_a is the operator from A to A defined for a given $a \in A$ as $U_a : x \mapsto 2a(ax) - a^2$ ($x \in A$). This operator is positive, i.e., $U_a(A_+) \subset A_+$. If $a \in \mathcal{L}(A)$, then $U_ax = a^2x$ ($x \in A$).

A weight (trace) Φ is called: *normal* if $\Phi(x) = \sup_\alpha \Phi(x_\alpha)$ for every increasing net (x_α) in A_+ with $x = \sup_\alpha x_\alpha$; *semifinite* if there exists an increasing net (a_α) in A_+ with $\sup_\alpha a_\alpha = \mathbb{1}$ and $\Phi(a_\alpha) \in \Lambda$ for all α ; *bounded* if $\Phi(\mathbb{1}) \in \Lambda$. Given two Λ -valued weights Φ and Ψ on A , say that Φ is dominated by Ψ if there exists $\lambda \in \Lambda_+$ such that $\Phi(x) \leq \lambda\Psi(x)$ for all $x \in A_+$.

4.9. We need a few additional remarks about descents and ascents. Fix $+\infty \in \mathbb{V}^{(\mathbb{B})}$. If $\Lambda = \mathcal{R}\downarrow$ and $\Lambda^\cup = \mathcal{R}\downarrow$ then

$$(\Lambda^\cup \cup \{+\infty\})\uparrow = (\Lambda \cup \{+\infty\})\uparrow = \Lambda\uparrow \cup \{+\infty\}\uparrow = \mathcal{R} \cup \{+\infty\}.$$

At the same time, $\Lambda^* := (\mathcal{R} \cup \{+\infty\})\downarrow = \text{mix}(\mathcal{R}\downarrow \cup \{+\infty\})$ consists of all elements of the form $\lambda_\pi := \text{mix}(\pi\lambda, \pi^\perp(+\infty))$ with $\lambda \in \Lambda^\cup$ and $\pi \in \mathbb{P}(\Lambda)$. Thus, $\Lambda^\cup \cup \{+\infty\}$ is a proper subset of Λ^* , since $x_\pi \in \Lambda \cup \{+\infty\}$ if and only if $\pi = 0$ or $\pi = I_\Lambda$.

Assume now that $A = \mathcal{A}\downarrow$ with \mathcal{A} a *JB*-algebra within $\mathbb{V}^{(\mathbb{B})}$ and \mathbb{B} equal to $\mathbb{P}(A)$. Every bounded weight $\Phi : A \rightarrow \Lambda$ is evidently extensional: $b := \llbracket x = y \rrbracket$ implies $bx = by$ which in turn yields $b\Phi(x) = \Phi(bx) = \Phi(by) = b\Phi(y)$ or, equivalently, $b \leq \llbracket \Phi(x) = \Phi(y) \rrbracket$. But an unbounded weight may fail to be extensional. Indeed, if $\Phi(x_0) = +\infty$ and $\Phi(x) \in \Lambda$ for some $x_0, x \in A$ and $b \in \mathbb{P}(A)$ then

$$\Phi(\text{mix}(bx, b^\perp x_0)) = \text{mix}(b\Phi(x), b^\perp(+\infty)) \notin \Lambda \cup \{+\infty\}.$$

Given a semifinite weight Φ on A , we define its extensional modification $\widehat{\Phi} : A \rightarrow \Lambda^*$ as follows: If $\Phi(x) \in \Lambda$ we put $\widehat{\Phi}(x) := \Phi(x)$. If $\Phi(x) = +\infty$ then $x = \sup D$ with $D := \{a \in A : 0 \leq a \leq x, \Phi(a) \in \Lambda\}$. Let b stand for the greatest element of $\mathbb{P}(\Lambda)$ such that $\Phi(bD)$ is order bounded in Λ^\cup and put $\lambda := \sup \Phi(bD)$. We define $\widehat{\Phi}(x)$ as $\lambda_b = \text{mix}(b\lambda, b^\perp(+\infty))$; i.e., $b\widehat{\Phi}(x) = \lambda$ and $b^\perp\widehat{\Phi}(x) = b^\perp(+\infty)$. It is not difficult to check that $\widehat{\Phi}$ is an extensional mapping. Thus, for $\varphi := \widehat{\Phi}\uparrow$ we have $\llbracket \varphi : \mathcal{A} \rightarrow \mathcal{R} \cup \{+\infty\} \rrbracket = \mathbb{1}$ and, according to [23, 1.6.6], $\widehat{\Phi} = \varphi\downarrow \neq \Phi$. But if we define $\varphi\downarrow$ as $\varphi\downarrow(x) = \varphi(x)$ whenever $\varphi(x) \in \Lambda$ and $\varphi\downarrow(x) = +\infty$ otherwise, then $\Phi = (\widehat{\Phi}\uparrow)\downarrow$.

4.10. Theorem. *Let A be a *JBW*-algebra and let τ be a normal semifinite real-valued trace on A . For each real-valued weight φ on A dominated by τ there exists a unique positive element $h \in A$ such that $\varphi(a) = \tau(U_{h^{1/2}}a)$ for all $a \in A_+$. Moreover, φ is bounded if and only if $\tau(h)$ is finite and φ is a trace if and only if h is a central element of A .*

◁ This fact was proved in [16]. ▷

4.11. Theorem. *Let A be a \mathbb{B} -*JBW*-algebra and let \mathbb{T} be a normal semifinite Λ -valued trace on A . For each weight Φ on A dominated by \mathbb{T} there exists a unique positive $h \in A$ such that $\Phi(x) = \mathbb{T}(U_{h^{1/2}}x)$ for all $x \in A_+$. Moreover, Φ is bounded if and only if $\mathbb{T}(h) \in \Lambda$ and Φ is a trace if and only if h is a central element of A .*

◁ We present a sketch of the proof. Taking into consideration the remarks in 4.9, we define $\varphi = \widehat{\Phi}\uparrow$ and $\psi = \widehat{\Psi}\uparrow$. Then within $\mathbb{V}^{(\mathbb{B})}$ the following hold: τ is a semifinite normal real-valued trace on \mathcal{A} and φ is real-valued weight on \mathcal{A} dominated by τ . By transfer we may apply Theorem 4.10 and find $h \in \mathcal{A}$ such that $\varphi(x) = \tau(U_{h^{1/2}}x)$ for all $x \in \mathcal{A}_+$. Actually, $h \in A$ and $\varphi\downarrow(x) = \tau\downarrow(U_{h^{1/2}}x)$ for all $x \in A_+$. It remains to note that $\Phi = \varphi\downarrow$ and $\mathbb{T} = \tau\downarrow$. The details of the proof are left to the reader. ▷

4.12. *JB*-algebras are nonassociative real analogs of C^* -algebras and von Neumann operator algebras. The theory of these algebras stems from Jordan, von Neumann, and Wigner [15] and exists as a branch of functional analysis since the mid 1960s. The stages of its development are reflected in Alfsen, Shultz, and Størmer [5]. The theory of *JB*-algebras undergoes intensive study, and the scope of its applications widens. Among the main areas of research are the structure and classification of *JB*-algebras, nonassociative integration and quantum probability theory, the geometry of states of *JB*-algebras, etc.; see Ajupov [1, 2]; Hanshe-Olsen and Størmer [14] as well as the references therein.

(2) The Boolean valued approach to *JB*-algebras was charted by Kusraev in the article [19] which contains Theorems 4.4 and 4.7 (also see [20]). These theorems are instances of the Boolean valued interpretation of the results by Shulz [34] and by Ajupov and Abdullaev [3]. In [19] Kusraev introduced the class of \mathbb{B} -*JBW*-algebras which is broader than the class of *JBW*-algebras. The principal distinction is that a \mathbb{B} -*JBW*-algebra has faithful representation as the algebra of selfadjoint operators in some AW^* -module rather than on a Hilbert space as in the case of *JBW*-algebras (cp. Kusraev and Kutateladze [22]). The class of *AJW*-algebras was firstly mentioned by Topping in [40]. Theorem 4.11 was not published before.

5. TRANSFER IN HARMONIC ANALYSIS

In what follows, G is a locally compact abelian group and τ is the topology of G , while $\tau(0)$ is a neighborhood base of 0 in G and G' stands for the dual group of G . Note that G is also the dual group of G' and we write $\langle x, \gamma \rangle := \gamma(x)$ ($x \in G$, $\gamma \in G'$).

5.1. By restricted transfer, G^\wedge is a group within $\mathbb{V}^{(\mathbb{B})}$. At the same time $\tau(0)^\wedge$ may fail to be a topology on G^\wedge . But G^\wedge becomes a topological group on taking $\tau(0)^\wedge$ as a neighborhood base of $0 := 0^\wedge$. This topological group is again denoted by G^\wedge itself. Clearly, G^\wedge may not be locally compact. Let U be a neighborhood of 0 such that U is compact. Then U is totally bounded. It follows by restricted transfer that U^\wedge is totally bounded as well, since total boundedness can be expressed by a restricted formula. Therefore the completion of G^\wedge is locally compact. The completion of G^\wedge is denoted by \mathcal{G} , and by the above observation \mathcal{G} is a locally compact abelian group within $\mathbb{V}^{(\mathbb{B})}$.

5.2. Let Y be a Dedekind complete vector lattice and let $Y_{\mathbb{C}}$ be the complexification of Y . A vector-function $\varphi : G \rightarrow Y$ is said to be *uniformly order continuous* on a set K if

$$\inf_{U \in \tau(0)} \sup\{|\varphi(g_1) - \varphi(g_2)| : g_1, g_2 \in K, g_1 - g_2 \in U\} = 0.$$

This amounts to saying that φ is order bounded on K , and if $e \in Y$ is an arbitrary upper bound of $\varphi(K)$, then for each $0 < \varepsilon \in \mathbb{R}$ there exists a partition of unity $(\pi_\alpha)_{\alpha \in \tau(0)}$ in $\mathbb{P}(Y)$ such that $\pi_\alpha |\varphi(g_1) - \varphi(g_2)| \leq \varepsilon e$ for all $\alpha \in \tau(0)$ and $g_1, g_2 \in K$, $g_1 - g_2 \in \alpha$. If, in this definition we put $g_2 = 0$, then we arrive at the definition of mapping *order continuous at zero*.

We now introduce the class of dominated Y -valued mappings. A mapping $\psi : G \rightarrow Y_{\mathbb{C}}$ is called *positive definite* if

$$\sum_{j,k=1}^n \psi(g_j - g_k) c_j \bar{c}_k \geq 0$$

for all finite collections $g_1, \dots, g_n \in G$ and $c_1, \dots, c_n \in \mathbb{C}$ ($n \in \mathbb{N}$).

For $n = 1$, the definition implies readily that $\psi(0) \in Y_+$. For $n = 2$, we have $|\psi(g)| \leq \psi(0)$ ($g \in G$). If we introduce the structure of an f -algebra with unit $\psi(0)$ in the order ideal of Y generated by $\psi(0)$ then, for $n = 3$, from the above definition we can deduce one more inequality

$$|\psi(g_1) - \psi(g_2)|^2 \leq 2\psi(0)(\psi(0) - \Re\psi(g_1 - g_2)) \quad (g_1, g_2 \in G).$$

It follows that every positive definite mapping $\psi : G \rightarrow Y_{\mathbb{C}}$ o -continuous at zero is order-bounded (by the element $\psi(0)$) and uniformly o -continuous. A mapping $\varphi : G \rightarrow Y$ is called *dominated* if there exists a positive definite mapping $\psi : G \rightarrow Y_{\mathbb{C}}$ such that

$$\left| \sum_{j,k=1}^n \varphi(g_j - g_k) c_j \bar{c}_k \right| \leq \sum_{j,k=1}^n \psi(g_j - g_k) c_j \bar{c}_k$$

for all $g_1, \dots, g_n \in G$, $c_1, \dots, c_n \in \mathbb{C}$ ($n \in \mathbb{N}$). In this case we also say that ψ is a *dominant* of φ . It can be easily shown that if $\varphi : G \rightarrow Y_{\mathbb{C}}$ has dominant order continuous at zero then φ is order bounded and uniformly order continuous.

We denote by $\mathfrak{D}(G, Y_{\mathbb{C}})$ the vector space of all dominated mappings from G into $Y_{\mathbb{C}}$ whose dominants are order continuous at zero. We also consider the set $\mathfrak{D}(G, Y_{\mathbb{C}})_+$ of all positive definite mappings from G into $Y_{\mathbb{C}}$. This set is a proper cone in $\mathfrak{D}(G, Y_{\mathbb{C}})$ and defines the order compatible with the structure of a vector space on $\mathfrak{D}(G, Y_{\mathbb{C}})$. Actually, $\mathfrak{D}(G, Y_{\mathbb{C}})$ is a Dedekind complete complex vector lattice; cp. 5.13 below. Also, define $\mathfrak{D}(\mathcal{G}, \mathcal{C}) \in \mathbb{V}^{(\mathbb{B})}$ to be the set of functions $\varphi : \mathcal{G} \rightarrow \mathcal{C}$ with the property that $\llbracket \varphi \text{ has dominant continuous at zero} \rrbracket = \mathbb{1}$.

5.3. Let $Y = \mathcal{R}\downarrow$. For every $\varphi \in \mathfrak{D}(G, Y_{\mathbb{C}})$ there exists a unique $\tilde{\varphi} \in \mathbb{V}^{(\mathbb{B})}$ such that $\llbracket \tilde{\varphi} \in \mathfrak{D}(\mathcal{G}, \mathcal{C}) \rrbracket = \mathbb{1}$ and $\llbracket \tilde{\varphi}(x^\wedge) = \varphi(x) \rrbracket = \mathbb{1}$ for all $x \in G$. The mapping $\varphi \mapsto \tilde{\varphi}$ is an linear and order isomorphism from $\mathfrak{D}(G, Y)$ onto $\mathfrak{D}(\mathcal{G}, \mathcal{C})\downarrow$.

5.4. Define $C_0(G)$ as the space of all continuous complex functions f on G vanishing at infinity. The latter means that for every $0 < \varepsilon \in \mathbb{R}$ there exists a compact set $K \subset G$ such that $|f(x)| < \varepsilon$ for all $x \in G \setminus K$. Denote by $C_c(G)$ the space of all continuous complex functions on G having compact support. Evidently, $C_c(G)$ is dense in $C_0(G)$ with respect to the norm $\|\cdot\|_\infty$.

5.5. Introduce the class of dominated operators. Let X be a complex normed space and let Y be a complex Banach lattice. A linear operator $T : X \rightarrow Y$ is said to be *dominated* or *having abstract norm* if T sends the unit ball of X into an order bounded subset of Y . This amounts to saying that there exists $c \in Y_+$ such that $|Tx| \leq c\|x\|_\infty$ for all $x \in C_0(Q)$. The set of all dominated operators from X to Y is denoted by $L_m(X, Y)$. If Y is Dedekind complete then

$$|T| := \{|Tx| : x \in X, \|x\| \leq 1\}$$

exists and is called the *abstract norm* or *dominant* of T . Moreover, if X is a vector lattice and Y is Dedekind complete then $L_m(X, Y)$ is a vector sublattice of $L^\sim(X, Y)$.

Given a positive $T \in L_m(C_0(G'), Y)$, we can define the mapping $\varphi : G \rightarrow Y$ by putting $\varphi(x) = T(\langle x, \cdot \rangle)$ for all $x \in G$, since $\gamma \mapsto \langle x, \gamma \rangle$ lies in $C_0(G')$ for every $x \in G$. It is not difficult to ensure that φ is order continuous at zero and positive definite. The converse is also true; see 5.8.

5.6. Consider a metric space (M, r) . The definition of metric space can be written as a bounded formula, say $\varphi(M, r, \mathbb{R})$, so that $\llbracket \varphi(M^\wedge, r^\wedge, \mathbb{R}^\wedge) \rrbracket = \mathbb{1}$ by restricted transfer. In

other words, (M^\wedge, r^\wedge) is a metric space within $\mathbb{V}^{(\mathbb{B})}$. Moreover we consider the internal function $r^\wedge : M^\wedge \rightarrow \mathbb{R}^\wedge \subset \mathcal{R}$ as an \mathcal{R} -valued metric on M^\wedge . Denote by (\mathcal{M}, ρ) the completion of (M^\wedge, r^\wedge) ; i.e., $\llbracket (\mathcal{M}, \rho) \text{ is a complete metric space} \rrbracket = \mathbb{1}$, $\llbracket M^\wedge \text{ is a dense subset of } \mathcal{M} \rrbracket = \mathbb{1}$, and $\llbracket r(x^\wedge) = \rho(x^\wedge) \rrbracket = \mathbb{1}$ for all $x \in M$.

Now, if $(X, \|\cdot\|)$ is a real (or complex) normed space then $\llbracket X^\wedge \text{ is a vector space over the field } \mathbb{R}^\wedge \text{ (or } \mathbb{C}^\wedge) \text{ and } \|\cdot\|^\wedge \text{ is a norm on } X^\wedge \text{ with values in } \mathbb{R}^\wedge \subset \mathcal{R} \rrbracket = \mathbb{1}$. So, we will consider X^\wedge as an \mathbb{R}^\wedge -vector space with \mathcal{R} -valued norm within $\mathbb{V}^{(\mathbb{B})}$. Let $\mathcal{X} \in \mathbb{V}^{(\mathbb{B})}$ stand for the (metric) completion of X^\wedge within $\mathbb{V}^{(\mathbb{B})}$. It is not difficult to see that $\llbracket \mathcal{X} \text{ is a real (complex) Banach space including } X^\wedge \text{ as an } \mathbb{R}^\wedge (\mathbb{C}^\wedge)\text{-linear subspace} \rrbracket = \mathbb{1}$, since the metric $(x, y) \mapsto \|x - y\|$ on X^\wedge is translation invariant. Clearly, if X is a real (complex) Banach lattice then $\llbracket \mathcal{X} \text{ is a real (complex) Banach lattice including } X^\wedge \text{ as an } \mathbb{R}^\wedge (\mathbb{C}^\wedge)\text{-linear sublattice} \rrbracket = \mathbb{1}$.

5.7. Theorem. *Let $Y = \mathcal{C}\downarrow$ and \mathcal{X}' be the topological dual of \mathcal{X} within $\mathbb{V}^{(\mathbb{B})}$. For every $T \in L_m(X, Y)$ there exists a unique $\tau \in \mathcal{X}'\downarrow$ such that $\llbracket \tau(x^\wedge) = T(x) \rrbracket = \mathbb{1}$ for all $x \in X$. The mapping $T \mapsto \phi(T) := \tau$ defines an isomorphism between the $\mathcal{C}\downarrow$ -modules $L_m(X, Y)$ and $\mathcal{X}'\downarrow$. Moreover, $|T| = |\phi(T)|$ for all $T \in L_m(X, Y)$. If X is a normed lattice then $\llbracket \mathcal{X}' \text{ is a Banach lattice} \rrbracket = \mathbb{1}$, while $\mathcal{X}'\downarrow$ is a vector lattice and ϕ is a lattice isomorphism.*

\triangleleft Suffice it to consider the real case. Apply [18, Theorem 8.3.2] to the lattice normed space $X := (X, |\cdot|)$ with $|x| = \|x\|\mathbb{1}$. By [18, Theorem 8.3.4 (1) and Proposition 8.3.4 (2)] the spaces $\mathcal{X}'\downarrow := \mathcal{L}^{(\mathbb{B})}(\mathcal{X}, \mathcal{R})\downarrow$ and $L_m(X, Y)$ are linear isometric. We are left with referring to [18, Proposition 5.5.1 (1)]. \triangleright

5.8. Theorem. *A mapping $\varphi : G \rightarrow Y_{\mathbb{C}}$ is order continuous at zero and positive definite if and only if there exists a unique positive operator $T \in L_m(C_0(G'), Y_{\mathbb{C}})$ such that $\varphi(x) = T(\langle x, \cdot \rangle)$ for all $(x \in G)$.*

\triangleleft By transfer, 5.3, and Theorem 5.7, we can replace φ and T by their Boolean valued representations $\tilde{\varphi}$ and τ . The norm completion of $C_0(G')^\wedge$ within $\mathbb{V}^{(\mathbb{B})}$ coincides with $C_0(\mathcal{G}')$. (This can be proved by the reasoning similar to that in Takeuti [37, Proposition 3.2].) Application of the classical Bochner Theorem (see Loomis [26, Section 36A]) to $\tilde{\varphi}$ and τ yields the desired result. \triangleright

5.9. We now specify the vector integral of use in this subsection; see details in [23, 5.14.B]. Let \mathcal{A} be a σ -algebra of subsets of Q , i.e. $\mathcal{A} \subset \mathcal{P}(Q)$. We identify this algebra with the isomorphic algebra of the characteristic functions $\{1_A := \chi_A : A \in \mathcal{A}\}$ so that $S(\mathcal{A})$ is the space of all \mathcal{A} -simple functions on Q ; i.e., $f \in S(\mathcal{A})$ means that $f = \sum_{k=1}^n \alpha_k \chi_{A_k}$ for some $\alpha_1, \dots, \alpha_n \in \mathbb{R}$ and disjoint $A_1, \dots, A_n \in \mathcal{A}$. Let a measure μ be defined on \mathcal{A} and take values in a Dedekind complete vector lattice Y . We suppose that μ is order bounded. If $f \in S(\mathcal{A})$ then we put

$$I_\mu := \int f d\mu = \sum_{k=1}^n \alpha_k \mu(A_k).$$

It can be easily seen that the integral I_μ can be extended to the spaces of μ -summable functions $\mathcal{L}^1(\mu)$ for which the more informative notations $\mathcal{L}^1(Q, \mu)$ and $\mathcal{L}^1(Q, \mathcal{A}, \mu)$ are also used. On identifying equivalent functions, we obtain the Dedekind σ -complete vector lattice $L^1(\mu) := L^1(Q, \mu) := L^1(Q, \mathcal{A}, \mu)$.

5.10. Assume now that Q is a topological space. Denote by $\mathcal{F}(Q)$, $\mathcal{K}(Q)$, and $\mathcal{B}(Q)$ the collections of all closed, compact, and Borel subsets of Q . A measure $\mu : \mathcal{B}(Q) \rightarrow Y$ is said to be *quasi-Radon* (*quasiregular*) if μ is order bounded and

$$\begin{aligned} |\mu|(U) &= \sup\{|\mu|(K) : K \in \mathcal{K}(Q), K \subset U\} \\ (|\mu|(U) &= \sup\{|\mu|(K) : K \in \mathcal{F}(Q), K \subset U\}). \end{aligned}$$

for every open (respectively, closed) set $U \subset Q$. If the above equalities are fulfilled for all Borel $U \subset Q$ then we speak about *Radon* and *regular* measures. Say that $\mu = \mu_1 + i\mu_2 : \mathcal{B}(Q) \rightarrow Y_{\mathbb{C}}$ have one of the above properties whenever this property is enjoyed by both μ_1 and μ_2 . We denote by $\text{qca}(Q, Y)$ the vector lattice of all σ -additive $Y_{\mathbb{C}}$ -valued quasi-Radon measures on $\mathcal{B}(Q)$. If Q is locally compact or even completely regular then $\text{qca}(Q, Y)$ is a vector lattice; see [18, Theorem 6.2.2]. The variation of a $Y_{\mathbb{C}}$ -valued (in particular, \mathbb{C} -valued) measure ν is denoted in the standard fashion: $|\nu|$.

5.11. Theorem. *Let Y be a real Dedekind complete vector lattice and let Q be a locally compact topological space. Then for each $T : L_m(C_0(Q), Y_{\mathbb{C}})$ there exists a unique measure $\mu := \mu_T \in \text{qca}(Q, Y_{\mathbb{C}})$ such that*

$$T(f) = \int_Q f d\mu \quad (f \in C_0(Q)).$$

The mapping $T \mapsto \mu_T$ is a lattice isomorphism from $L_m(C_0(Q), Y_{\mathbb{C}})$ onto $\text{qca}(Q, Y_{\mathbb{C}})$.

◁ See [24, Theorem 2.5]. ▷

5.12. Theorem. *Let G be a locally compact abelian group, let G' be the dual group of G , and let Y be a Dedekind complete real vector lattice. For a mapping $\varphi : G \rightarrow Y_{\mathbb{C}}$ the following are equivalent:*

- (1) φ has dominant order continuous at zero.
- (2) There exists a unique measure $\mu \in \text{qca}(G', Y_{\mathbb{C}})$ such that

$$\varphi(g) = \int_{G'} \chi(g) d\mu(\chi) \quad (g \in G).$$

◁ This is immediate from Theorems 5.8 and 5.11. ▷

5.13. Corollary. *The Fourier transform establishes an order and linear isomorphism between the space of measures $\text{qca}(G', Y)$ and the space of dominated mappings $\mathfrak{D}(G, Y_{\mathbb{C}})$. In particular, $\mathfrak{D}(G, Y_{\mathbb{C}})$ is a Dedekind complete complex vector lattice.*

5.14. (1) In [37] Takeuti introduced the Fourier transform for the mappings defined on a locally compact abelian group and having as values pairwise commuting normal operators in a Hilbert space. By applying the transfer principle, he developed a general technique for translating the classical results to operator-valued functions. In particular, he established a version of the Bochner Theorem describing the set of all inverse Fourier transforms of positive operator-valued Radon measures. Given a complete Boolean algebra \mathbb{B} of projections in a Hilbert space H , denote by (\mathbb{B}) the space of all selfadjoint operators on H whose spectral resolutions are in \mathbb{B} ; i.e., $A \in (\mathbb{B})$ if and only if $A = \int_{\mathbb{R}} \lambda dE_{\lambda}$ and $E_{\lambda} \in \mathbb{B}$ for all $\lambda \in \mathbb{R}$. If $Y := (\mathbb{B})$ then Theorem 5.8 is essentially Takeuti's result [37, Theorem 1.3].

(2) Kusraev and Malyugin in [24] abstracted Takeuti's results in the following directions: First, they considered more general arrival spaces, namely, norm complete lattice normed spaces. So the important particular cases of Banach spaces and Dedekind complete vector lattices were covered. Second, the class of dominated mappings was identified with the set of all inverse Fourier transforms of order bounded quasi-Radon vector measures. Third, the construction of a suitable Boolean valued universe was eliminated from all definitions and statements of results.

In particular, Theorem 5.12 and Corollary 5.13 correspond to [24, Theorem 4.3] and [24, Theorem 4.4]; while their lattice normed versions, to [24, Theorem 4.1] and [24, Theorem 4.5], respectively.

(3) Theorem 5.7 is due to Gordon [11, Theorem 2]. Proposition 3.3 in Takeuti [37] is essentially the same result for the particular departure and arrival spaces; i.e., $X = L^1(G)$ and $Y = (\mathbb{B})$. Theorem 5.11 is taken from Kusraev and Malyugin [24]. In the case of Q compact, it was proved by Wright in [44, Theorem 4.1]. In this result μ cannot be chosen regular rather than quasiregular. The quasiregular measures were introduced by Wright in [43].

REFERENCES

- [1] *Ajupov Sh. A.* Jordan Operator Algebras // J. Soviet Math.—1987.—Vol. 37, No. 6.—P. 1422–1448.
- [2] *Ajupov Sh. A.* Classification and Representation of Ordered Jordan Algebras.—Tashkent: Fan, 1986.—121 p. [in Russian]
- [3] *Ajupov Sh. A. and Abdullaev R. Z.*, Radon–Nikodým theorem for weights on semi-finite JBW-algebras // Math. Z.—1985.—Vol. 188.—P. 475–484.
- [4] *Albeverio S., Ajupov Sh. A., and Kudaybergenov K. K.* Derivations on the algebras of measurable operators affiliated with a type I von Neumann algebra // Siberian Adv. Math.—2008.—Vol. 18, No. 2.—P. 86–94.
- [5] *Alfsen E. M., Shultz F. W., and Störmer E.* A Gelfand–Neumark theorem for Jordan algebras // Adv. Math.—1978.—Vol. 28, No. 1.—P. 11–56.
- [6] *Aliprantis C. D. and Burkinshaw O.* Positive Operators.—London etc.: Acad. Press Inc., 1985.—xvi+367 p.
- [7] *Bell J. L.* Set Theory: Boolean-Valued Models and Independence Proofs in Set Theory.—Oxford: Clarendon Press, 2005.—xxii+191 p.—(Oxford Logic Guides 47).
- [8] *Ber A. F., de Pagter B., and Sukochev F. A.* Derivations in algebras of operator-valued functions // J. Operator Theory.—2011.—Vol. 66, No. 2.—P. 261–300.
- [9] *Berberian S. K.* Baer *-Rings.—Berlin: Springer-Verlag, 1972 (Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen. Band 195).—296 p.
- [10] *Chilin V. I.* Partially ordered Baer involutive algebras // J. Soviet Math.—1987.—Vol. 37, No. 6.—P. 1449–1472.
- [11] *Gordon E. I.* K -spaces in Boolean-valued models of set theory // Sov. Math., Dokl.—1981.—Vol. 23.—P. 579–582.
- [12] *Gutman A. E.* Locally one-dimensional K -spaces and σ -distributive Boolean algebras // Siberian Adv. Math.—1995.—Vol. 5, No. 2.—P. 99–121.
- [13] *Gutman A. E., Kusraev A. G., and Kutateladze S. S.* The Wickstead Problem // Siberian Elektron. Math. Reports.—2008.—Vol. 5.—P. 293–333.
- [14] *Hanshe-Olsen H. and Störmer E.* Jordan Operator Algebras.—Boston: Pitman Publ. Inc., 1984.—183 p.
- [15] *Jordan PP., von Neumann J., and Wigner E.* On an algebraic generalization of the quantum mechanic formalism // Ann. Math.—1944.—Vol. 35.—P. 29–64.
- [16] *King W. PP. C.* Semi-finite traces on JBW-algebras // Math. Proc. Cambridge Philos. Soc.—1983.—Vol. 72.—P. 323–326.
- [17] *Korol' A. M. and Chilin V. I.*, Measurable operators in a Boolean-valued model of set theory // Dokl. Akad. Nauk UzSSR.—1989.—Vol. 3.—P. 7–9. [in Russian]

- [18] *Kusraev A. G.* Dominated Operators.—Kluwer: Dordrecht, 2000.—xiii+446 p.
- [19] *Kusraev A. G.* Boolean-valued analysis and JB -algebras // *Siberian Math. J.*—1994.—Vol. 35, No. 1.—P. 114–122.
- [20] *Kusraev A. G.* On the structure of type I_2 AJW -algebras // *Siberian Math. J.*—1999.—Vol. 40, No. 4.—P. 764–774.
- [21] *Kusraev A. G. and Kutateladze S. S.* Boolean Valued Analysis.—Dordrecht: Kluwer Academic Publishers, 1999.—xii+322 p.
- [22] *Kusraev A. G. and Kutateladze S. S.* Introduction to Boolean Valued Analysis.—M.: Nauka, 2005.—526 p. [in Russian]
- [23] *Kusraev A. G. and Kutateladze S. S.* Boolean Valued Analysis: Selected Topics.—Vladikavkaz: SMI VSC RAS, 2014.—iv+400 p.—(Trends in Science: The South of Russia. A Math. Monogr. Issue 6).
- [24] *Kusraev A. G. and Malugin S. A.* On the Fourier transform of dominated mappings // *Siberian Math. J.*—1994.—Vol. 35, No. 6.—P. 1141–1156.
- [25] *Kutateladze S. S.* What is Boolean valued analysis? // *Siberian Adv. Math.*—2007.—Vol. 17, No. 2.—P. 91–111.
- [26] *Loomis L. H.* An Introduction to Abstract Harmonic Analysis.—Princeton: D. Van Nostrand Company Inc., 1953.—vii+190 p.
- [27] *Luxemburg W. A. J. and Zaanen A. C.* Riesz Spaces. Vol. 1.—Amsterdam–London: North-Holland, 1971.—514 p.
- [28] *Maslyuchenko O. V., Mykhaylyuk V. V., and Popov M. M.* A lattice approach to narrow operators // Positivity.—2009.—Vol. 13.—P. 459–495.
- [29] *Plichko A. M. and Popov M. M.* Symmetric function spaces on atomless probability spaces. *Dissertationes Math. (Rozprawy Mat.)*.—1990.—Vol. 306.—P. 1–85.
- [30] *Popov M. M. and Randrianantoanina B.* Narrow Operators on Function Spaces and Vector Lattices.—Berlin: De Gruyter, 2012.—xii+319 p. (De Gruyter Stud. in Math., Vol. 45).
- [31] *Rosenthal H. P.* Embeddings of L^1 in L^1 // *Contemp. Math.*—1984.—Vol. 26.—P. 335–349.
- [32] *Sakai S.* C^* -Algebras and W^* -Algebras.—Berlin etc.: Springer-Verlag, 1971.—ix+256 p.
- [33] *Segal I. E.* A non-commutative extension of abstract integration // *Ann. of Math.*—1953.—Vol. 57.—P. 401–457.
- [34] *Shultz F. W.* On normed Jordan algebras which are Banach dual spaces // *J. Funct. Anal.*—1979.—Vol. 31, No. 3.—P. 360–376.
- [35] *Tabuev S. N.* Decomposition of an atomic operator // *Vladikavkaz Math. J.*—2003.—Vol. 5, No. 1.—P. 53–56.
- [36] *Takeuti G.* Two Applications of Logic to Mathematics.—Tokyo and Princeton: Iwanami and Princeton Univ. Press, 1978.—137 p.
- [37] *Takeuti G.* A transfer principle in harmonic analysis // *J. Symbolic Logic.*—1979.—Vol. 44, No. 3.—P. 417–440.
- [38] *Takeuti G.* Boolean valued analysis // In: Applications of Sheaves (Proc. Res. Sympos. Appl. Sheaf Theory to Logic, Algebra and Anal., Univ. Durham, Durham, 1977).—Berlin etc.: Springer-Verlag, 1979.—P. 714–731.—(Lecture Notes in Math., 753).
- [39] *Takeuti G. and Zaring W. M.* Axiomatic set Theory.—N. Y.: Springer-Verlag, 1973.—238 p.
- [40] *Topping D. M.* Jordan algebras of self-adjoint operators // *Mem. Amer. Math. Soc.*—1965.—Vol. 53.—48 p.
- [41] *Vulikh B. Z.* Introduction to the Theory of Partially Ordered Spaces.—Groningen: Wolters-Noordhoff, 1967.—xv+387 p.
- [42] *Wickstead A. W.* Representation and duality of multiplication operators on Archimedean Riesz spaces // *Comp. Math.*—1977.—Vol. 35, No. 3.—P. 225–238.
- [43] *Wright J. D. M.* Stone algebra valued measures and integrals // *Proc. London Math. Soc.*—1969.—Vol. 19, No. 3.—P. 107–122.
- [44] *Wright J. D. M.* An extension theorem // *J. London Math. Soc.*—1973.—Vol. 7, No. 2.—P. 531–539.
- [45] *Zaanen A. C.* Riesz Spaces. Vol. 2.—Amsterdam etc.: North-Holland, 1983.—xi+720 p.

Anatoly G. Kusraev and Semën S. Kutateladze

SOME ASPECTS OF BOOLEAN VALUED ANALYSIS

Препринт № 294

Ответственный за выпуск
А. Е. Гутман

Издание подготовлено с использованием макропакета $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{T}\mathcal{E}\mathcal{X}$,
разработанного Американским математическим обществом

This publication was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{T}\mathcal{E}\mathcal{X}$,
the American Mathematical Society's $\text{T}\mathcal{E}\mathcal{X}$ macro package

Подписано в печать 14.02.15. Формат 60×84 ψ s.
Усл. печ. л. 2,0. Уч.-изд. л. 1,5. Тираж 75 экз. Заказ № 5.

Отпечатано в ООО «Омега Принт»
пр. Академика Лаврентьева, 6, 630090 Новосибирск