

INVARIANTS OF VIRTUAL LINKS

YULIYA MIKHALCHISHINA

The virtual knot theory has been introduced by Kauffman[1] as a generalization of the classical knot theory. Virtual knots (and links) are represented as generic immersions of circles in the plane (virtual link diagrams) where double points can be classical (with the usual information on overpasses and underpasses) or virtual. Virtual link diagrams are equivalent under ambient isotopy and some types of local moves (generalized Reidemeister moves).

Using virtual generalized Reidemeister moves we can introduce a notion of "virtual" braids. Virtual braids on n strands form a group denoted by VB_n . The relation between virtual braids and virtual knots (and links) are completely determined by a generalization of Alexander and Markov Theorem [2, 3]. It is worth to mention that for virtual braids an Alexander-like theorem states that any virtual link can be represented as the closure of a virtual braid.

In the classical case it is known that the braid group embeds into $\text{Aut}(F_n)$ by Artin representation which is a local one. Wada [6] constructed local representations of the braid group B_n into $\text{Aut}(F_n)$. There are three types. It is proved [7] that these representations are faithful.

Proposition. *For every Wada representation*

$$w_1^r, w_2, w_3 : B_n \rightarrow \text{Aut}F_n, \quad r \in \mathbb{Z}$$

it is possible to construct the corresponding representation of the virtual braid group

$$W_1^r, W_2, W_3 : VB_n \rightarrow \text{Aut}F_{n+1}, \quad r \in \mathbb{Z},$$

such that the restriction each of them onto B_n is coincide with the corresponding Wada representation, i.e.

$$W_k|_{B_n} = w_k, \quad k = 1, 2, 3.$$

Analogously to the way shown in [4] we introduce the notion of the group of the virtual link $G(L_v)$ for representations of the virtual braid group W_k , $k = 1, 2, 3$. Let the virtual link $L_v = \widehat{\beta}_v$ be a closure of the virtual braid, where $\beta_v \in VB_n$. We define

$$G_k(L_v) = \langle x_1, x_2, \dots, x_n, y \mid x_i = W_k(\beta_v)(x_i), \quad i = 1, 2, \dots, n \rangle, \quad k = 1, 2, 3.$$

Theorem. *The constructed groups $G_k(L_v)$, $k = 1, 2, 3$, are invariants of the virtual link L_v .*

REFERENCES

- [1] L. Kauffman, "Virtual knot theory", *Eur. J. Comb.*, Vol. 20, No. 7, 663-690, 1999.
- [2] S. Kamada, "Invariants of virtual braids and a remark on left stabilisations and virtual exchange moves", *Kobe J. Math.*, Vol. 21, 33-49, 2004.
- [3] L. Kauffman, S. Lambropoulou, "The LMove and Virtual Braids", *JKTR*, Vol. 15, No. 6, 773-811, 2006.
- [4] V. Bardakov, P. Bellingeri, "Groups of virtual and welded links", *J. Knot Theory Ramifications*, Vol. 23, No. 3, 23, 2014.
- [5] J. Birman, *Braid, links and mapping class group*, Princeton Univ. press, 1975.
- [6] M. Wada, "Group invariants of links", *Topology*, Vol. 31, 399-406, 1992.
- [7] V. Shpilrain, "Representing braids by automorphisms", *Int.J. Algebra Comput.*, Vol. 11, No. 6, 773-777, 2001.

SOBOLEV INSTITUTE OF MATHEMATICS, ACAD. KOPTYUG AVENUE, 4, NOVOSIBIRSK, 630090, RUSSIA
E-mail address: mikhailchishina@math.nsc.ru

The research was supported by the Russian-Chinese grant (14-01-91161).