

Operads of decorated trees

Pavel Kolesnikov
Sobolev Institute of Mathematics
Novosibirsk, Russia

Mal'cev Meeting-2013, Novosibirsk
November 11-15, 2013

Algebras

Associative, alternative,
Lie, Jordan, Malcev, Poisson,
Lie&Jordan triple systems,
Sabinin algebras, differential algebras,
Hom-algebras,

Splitting



Replicaton



pre-, post-Algebras
(Dendriform algebras)

Koszul duality



di-, tri-Algebras
(Loday-type algebras)

Algebras

Associative, alternative,
Lie, Jordan, Malcev, Poisson,
Lie&Jordan triple systems,
Sabinin algebras, differential algebras,
Hom-algebras,

Splitting

Replication

pre-, post-Algebras
(Dendriform algebras)

Koszul duality



di-, tri-Algebras
(Loday-type algebras)

Example:

Lie Algebras

$$xy + yx = 0$$

$$(xy)z + (yz)x + (zx)y = 0$$



Left-symmetric algebras
(pre-Lie algebras)

$$(xy)z - x(yz) = (yx)z - y(xz)$$

[E. Vinberg, 1960]

[J.-L. Koszul, 1961]

[M. Gerstenhaber, 1963]


[A. Cayley, 1896]

Leibniz Algebras

$$x(yz) = (xy)z + y(xz)$$

[A. Bloh, 1965]

[J.-L. Loday, 1993]



2d-Conformal field theory

V space of states

$T : V \rightarrow V$ translation operator

$Y : V \rightarrow \text{gl } V[[z, z^{-1}]]$ state-field correspondence

$$a \mapsto Y(a, z) = \sum_{n \in \mathbb{Z}} a(n) z^{-n-1}$$

+ axioms [R. Borcherds, 1986], [V. Kac, 1996]

Locality: $[Y(a, z), Y(b, w)](z - w)^N = 0$ for $N \gg 0$

T-invariance: $[T, Y(a, z)] = \partial_z Y(a, z)$

$Y(a, z)Y(b, z) = \text{undefined}$

$Y(a, z)Y(b, w)$ has a singularity in $z = w$

$Y(a, z)Y(b, z) = \text{undefined}$

$Y(a, z)Y(b, w)$ has a singularity in $z = w$

Operator product expansion formula:


$$Y(a, z)Y(b, w) = \sum_{n=0}^{N(a,b)} \frac{Y(c_n, w)}{(z-w)^{n+1}} + \langle \text{principal part} \rangle$$



$$c_n = [a_{(n)}b], n \geq 0:$$

Algebraic operations on V

$(V, [\cdot_{(0)}\cdot])$ Leibniz algebra



$$= Y(:ab:, w) + (z-w)(\dots)$$

Wick product

$(V, : \cdot \cdot :)$ pre-Lie algebra

[M. Rosellen, 2005]

Classical Yang-Baxter equation (dynamic)

$$X_{13}(u+v)X_{23}(v) - X_{23}(v)X_{12}(u) - X_{21}(u)X_{13}(u+v) = 0$$

A Algebra

$X : \mathcal{D} \rightarrow A \otimes A$ meromorphic function

$$\mathcal{D} \subset \mathbb{C}$$

$$X_{23} = 1 \otimes X$$

$$X_{21} = X^{(12)} \otimes 1, \text{ etc.}$$

If A has nondegenerate symmetric associative bilinear form $\langle \cdot, \cdot \rangle$

then

$$X(u) \in A \otimes A \quad \Leftrightarrow \quad P_u(\cdot) : A \rightarrow A$$

$$X = \sum x_i \otimes y_i \quad \Leftrightarrow \quad P_u(x) = \sum \langle x_i, x \rangle y_i$$

CYBE (operator form):

$$P_{u+v}(x)P_v(y) = P_v(P_u(x)y) + P_{u+v}(xP_u(y))$$

For Lie algebras [A. Belavin, V. Dinfeld, 1982]

$$[P_{u+v}(x), P_v(y)] = P_v([P_u(x), y]) + P_{u+v}([x, P_u(y)])$$

$X \equiv \text{const}$:

$$[P(x), P(y)] = P([P(x), y] + [x, P(y)]) \quad P : L \rightarrow L \text{ is a Baxter operator}$$

[M. Semenov-Tian-Shansky, 1983]

$x * y = [P(x), y], x, y \in L$: left-symmetric product

$(L, *)$ pre-Lie algebra

For Lie algebras [A. Belavin, V. Dinfeld, 1982]

$$[P_{u+v}(x), P_v(y)] = P_v([P_u(x), y]) + P_{u+v}([x, P_u(y)])$$

$X \equiv \text{const}$:

$$[P(x), P(y)] = P([P(x), y] + [x, P(y)]) \quad P : L \rightarrow L \text{ is a Baxter operator}$$

[M. Semenov-Tian-Shansky, 1983]

$x * y = [P(x), y], x, y \in L$: left-symmetric product

$(L, *)$ pre-Lie algebra

Define $x_\lambda = \text{Res}_{u=0} e^{\lambda u} P_u(x) \in L[\lambda]$

Then

$$[x_\lambda, y_\mu] = [x_\lambda, y]_{\lambda+\mu}$$

Conformal Leibniz identity for $[x_{(n)}y]$ given by $[x_\lambda, y] = \sum_{n \geq 0} \frac{\lambda^n}{n!} [x_{(n)}y]$

Operad:

Collection of linear spaces $\mathcal{C}(n)$, $n \geq 1$, with

Composition rule $\mathcal{C}(n) \otimes \mathcal{C}(m_1) \otimes \cdots \otimes \mathcal{C}(m_n) \rightarrow \mathcal{C}(m_1 + \cdots + m_n)$

Symmetric group action $S_n \curvearrowright \mathcal{C}(n)$

Identity map $I \in \mathcal{C}(1)$


Var variety of algebras

Var $\langle X \rangle$ free algebra in Var generated by $X = \{x_1, x_2, \dots\}$


Polylinear part of Var $\langle X \rangle$ is an operad

Var (n) polynomials of degree n in x_1, \dots, x_n


Homological algebra,
Geometry, Math.Physics, ...



Homological algebra,
Geometry, Math.Physics, ...



Standard n-simplices




Free operad:

$$\Omega = \bigcup_{n \geq 1} \Omega_n$$

$\mathcal{F}(N)$ spanned by all planar trees with N leaves, vertices labelled by $f \in \Omega$

Order of a vertex = n if $f \in \Omega_n$




Free decorated operad: $\Omega = \bigcup_{n \geq 1} \Omega_n$

$\mathcal{F}(N)$ spanned by all decorated planar trees with N leaves

Composition is defined in such a way that:

- (1) Attaching a tree to a non-decorated leaf removes decoration
- (2) Attaching a tree to a decorated leaf preserves decoration



Replicating an operad

Perm:

$$x(yz) = (xy)z$$

$$(xy)z = (yx)z$$

$$\dim \text{Perm}(n) = n:$$

$$e_i^{(n)} = (x_1 \overset{i}{\wedge} \dots \wedge x_n) x_i$$

ComTrias: Perm &

$$x \perp y = y \perp x$$

$$(x \perp y) \perp z = x \perp (y \perp z)$$

$$(x \perp y)z = (xy)z$$

$$x(y \perp z) = (xy) \perp z$$

$$\dim \text{ComTrias}(n) = 2^n - 1:$$

$$e_{i_1, \dots, i_k}^{(n)} = (x_1 \overset{i_1}{\wedge} \dots \wedge \dots \wedge \overset{i_k}{\wedge} \dots \wedge x_n) (x_{i_1} \perp \dots \perp x_{i_k})$$

Replicating an operad

Example:

(C, \cdot) commutative algebra over a field \mathbb{k}

$\varepsilon : C \rightarrow \mathbb{k}$ homomorphism

e.g., counit of a bialgebra

Then

$$xy = \varepsilon(x) \cdot y$$

$$x \perp y = x \cdot y$$

turns C into a ComTrias-algebra

$C = \mathbb{k}\Gamma$ group algebra

$$\varepsilon(g) = 1, g \in \Gamma$$

Replicating an operad

[F. Chapoton, 2001]: $\text{Leib} = \text{Lie} \circ \text{Perm}$ (Manin white product of operads
[V. Ginzburg, M. Kapranov, 1994])

[B. Vallette, 2008]: $\mathcal{P} \circ \text{Perm} = \mathcal{P} \otimes \text{Perm}$ (Hadamard product of operads)
 $\mathcal{P} \circ \text{ComTrias} = \mathcal{P} \otimes \text{ComTrias}$

for binary quadratic operads

Definition:

$$\text{di-}\mathcal{P} = \mathcal{P} \otimes \text{Perm} \quad \text{tri-}\mathcal{P} = \mathcal{P} \otimes \text{ComTrias}$$

[P.K., 2008]
[A. Pozhidaev, 2009] } binary case (di-)

[M. Bremner, R. Felipe, J. Sanchez-Ortega, 2011] - general case (di-)
[V. Gubarev, P.K., 2011] - binary case (tri-)

[J.PeI, C. Bai, L. Guo, X. Ni, 2012] - name of the procedure

Replicating an operad
(general tri-algebra case)

$$\Omega = \{f_1, f_2, \dots\}$$

$$\nu(f_k) = n_k \geq 1$$

\mathcal{P} -Algebra:

$$(A, \Omega^A)$$

$$f_k^A : \underbrace{A \otimes \dots \otimes A}_{n_k} \rightarrow A$$

Defining identities Σ (multi-linear)

$$\Omega_3 = \{f_1^H, f_2^H, \dots\}$$

$$\nu(f_k^H) = n_k \geq 1$$

$$H \subseteq \{1, \dots, n_k\}, H \neq \emptyset$$

tri- \mathcal{P} -Algebra:

$$(D, \Omega_3^D)$$

Defining identities Σ_3 (multi-linear)



$$C = \text{ComTrias}\langle x_1, x_2, \dots \rangle$$

$$\boxed{A \models \Sigma \iff C \otimes A \models \Sigma_3}$$

$$f^H(c_1 \otimes a_1, \dots, c_n \otimes a_n) = e_H^{(n)}(c_1, \dots, c_n) \otimes f(a_1, \dots, a_n)$$

Identities Σ_3 may be derived explicitly from Σ

Bremner & Sanchez-Ortega + Gubarev & K.
(Perm instead of ComTrias)

Replicating an operad

Examples

di-Lie = Leib

$$\mu(x, y) = -\mu(y, x) \Rightarrow \boxed{\mu^1(x, y) = -\mu^2(y, x)}$$

$xy = \mu^2(x, y) \Rightarrow$ Left Leibniz algebra

$xy = \mu^1(x, y) \Rightarrow$ Right Leibniz algebra

di-Com = Perm [F. Chapoton, 2001]

di-As = Dias [J.-L. Loday, T. Pirashvili, 1993]

di-Alt [L. Dong, 2005]

di-Jord [M. Bremner, L. Peresi, 2011]

di-JTS [M. Bremner, R. Felipe, J. Sanchez-Ortega, 2011]

di-Mal [M. Bremner, L. Peresi, J. Sanchez-Ortega, 2011]

tri-Com = ComTrias [B. Vallette, 2007]

tri-As = Trias [J.-L. Loday, M. Ronco, 2004]

Replicating an operad

Theorem

Let $\nu(f) \geq 2$ for all $f \in \Omega$.

Then for every (di-)tri- \mathcal{P} -algebra D there exists canonical \mathcal{P} -algebra \widehat{D} and a (Perm-) ComTrias-algebra C such that

$$D \subset C \otimes \widehat{D}$$

Remark.

Theorem remains valid for algebras with derivations or endomorphisms in the language.

Corollary.

$$\dim \text{di-}\mathcal{P}(n) = n \dim \mathcal{P}(n)$$

$$\dim \text{tri-}\mathcal{P}(n) = (2^n - 1) \dim \mathcal{P}(n)$$

Replicating an operad (PBW-type problems)

$\omega : \mathcal{R} \rightarrow \mathcal{P}$ morphism of operads

functor $\omega : \mathcal{P}$ -algebras \rightarrow \mathcal{R} -algebras

$$A \mapsto A^{(\omega)}$$

e.g.,

As \rightarrow Lie

$$A \mapsto A^{(-)}, [x, y] = xy - yx$$

As \rightarrow Jord

$$A \mapsto A^{(+)}, x \cdot y = xy + yx$$

? Whether every \mathcal{R} -algebra L may be embedded into $A^{(\omega)}$
for an appropriate \mathcal{P} -algebra A ?

(Embedding problem)

? If $\dim L < \infty$, whether minimal $\dim A$ is finite?

(Ado problem)

? What is the structure of the universal enveloping \mathcal{P} -algebra $U(L)$
of a given \mathcal{R} -algebra L ?

(Poincaré-Birkhoff-Witt problem)

Replicating an operad

(PBW-type problems)

A morphism $\omega : \mathcal{R} \rightarrow \mathcal{P}$

induces

$$\omega \otimes \text{id} : \mathcal{R} \otimes \mathcal{C} \rightarrow \mathcal{P} \otimes \mathcal{C}$$

where $\mathcal{C} = \text{Perm}$ or ComTrias

Hence, we have natural functors

$\omega \otimes \text{id} : \text{di-}\mathcal{P}\text{-algebras} \rightarrow \text{di-}\mathcal{R}\text{-algebras}$

$\text{tri-}\mathcal{P}\text{-algebras} \rightarrow \text{tri-}\mathcal{R}\text{-algebras}$

Replicating an operad (PBW-type problems)

Example:

$$[x, y] = xy - yx$$

$$\gamma(x, y) = \mu(x, y) - \mu^{(12)}(x, y) \mid \otimes e_1^{(2)} \in \text{Perm}(2)$$

$$\gamma(x, y) \otimes e_1^{(2)} = \mu(x, y) \otimes e_1^{(2)} - \mu^{(12)}(x, y) \otimes e_1^{(2)}$$

$$\begin{array}{ccccc}
 \parallel & & \parallel & & \parallel \\
 \gamma^1(x, y) & & \mu^1(x, y) & & (\mu(x, y) \otimes e_2^{(2)})^{(12)} = \mu^2(x, y)^{(12)} \\
 \parallel & & \parallel & & \parallel \\
 [x \dashv y] & & x \dashv y & & y \vdash x
 \end{array}$$

$$[x \dashv y] = x \dashv y - y \vdash x$$

Replicating an operad
(PBW-type problems)

The same problems make sense in these settings

Replicating an operad
(PBW-type problems)

The same problems make sense in these settings

$$(C \otimes A)^{(\omega \otimes \text{id})} \simeq C \otimes A^{(\omega)}$$

$C \in \text{Perm or ComTrias}$

Corollary.

(1) If the Embedding problem has positive solution for $(\mathcal{R}, \mathcal{P}, \omega)$, then it has positive solution for $(\text{di-}\mathcal{R}, \text{di-}\mathcal{P}, \omega \otimes \text{id})$ and $(\text{tri-}\mathcal{R}, \text{tri-}\mathcal{P}, \omega \otimes \text{id})$;

(2) If the Ado problem has positive solution for $(\mathcal{R}, \mathcal{P}, \omega)$, then it has positive solution for $(\text{di-}\mathcal{R}, \text{di-}\mathcal{P}, \omega \otimes \text{id})$ and $(\text{tri-}\mathcal{R}, \text{tri-}\mathcal{P}, \omega \otimes \text{id})$;

(3) For a (di-)tri- \mathcal{R} -algebra L , the universal enveloping (di-)tri- \mathcal{P} -algebra $U(L)$ is isomorphic to the subalgebra of $C \otimes U(\widehat{L})$ generated by the image of L .

e.g., $C = \mathbb{k}\mathbb{Z}_2$

Replicating an operad (Special identities)

$\omega : \mathcal{R} \rightarrow \mathcal{P}$ morphism of operads

functor $\omega : \mathcal{P}\text{-algebras} \rightarrow \mathcal{R}\text{-algebras}$

$$A \mapsto A^{(\omega)}$$

\mathcal{R} -algebra L is *special* if $L \subseteq A^{(\omega)}$

for some \mathcal{P} -algebra A .

$S\mathcal{R}$ = operad, governing the variety generated by all special \mathcal{R} -algebras

$$\mathcal{R} \rightarrow S\mathcal{R}$$

Kernel = *Special identities*

Replicating an operad (Special identities)

Theorem (V. Gubarev, P.K., V. Voronin)

For any $\omega : \mathcal{R} \rightarrow \mathcal{P}$, consider

$$\omega \otimes \text{id} : \text{di-}\mathcal{R} \rightarrow \text{di-}\mathcal{P}$$

$$\omega \otimes \text{id} : \text{tri-}\mathcal{R} \rightarrow \text{tri-}\mathcal{P}$$

Then

$$S\text{di-}\mathcal{R} = \text{di-}S\mathcal{R}, \quad S\text{tri-}\mathcal{R} = \text{tri-}S\mathcal{R}.$$

Corollary.

- (1) For $\text{di-As} \rightarrow \text{di-Jord}$: No special identities of degree ≤ 7
[M. Bremner, L. Perezi, 2011]
- (2) Special identities for $\text{di-As} \rightarrow \text{di-JTS}$
[M. Bremner, R. Felipe, J. Sanchez-Ortega, 2011]
- (3) Special identities for $\text{di-Lie} = \text{Leib} \rightarrow \text{di-LTS}$
[M. Bremner, J. Sanchez-Ortega, 2011]
- (4) For $\text{di-Alt} \rightarrow \text{di-Mal}$: No special identities of degree < 7
[M. Bremner, L. Perezi, J. Sanchez-Ortega, 2011]

Replicating an operad (Mal'cev conjugacy)

Levi–Mal'cev Theorem

L Lie algebra ($L \in \text{di-Lie}$)

R max solvable ideal

$L = S \ltimes R$, $S \simeq L/R$ semisimple Lie algebra

All such S are conjugate by inner automorphisms

J. Mason (2013):

L Leibniz algebra ($L \in \text{di-Lie}$)

R max solvable ideal

$L = S \ltimes R$, $S \simeq L/R$ semisimple Lie algebra

Such S may not be conjugate by inner automorphism

Splitting an operad

Algebra $(A, *)$

(A, \prec, \succ) : $x * y = x \prec y + x \succ y$ (pre-algebra)

(A, \prec, \succ, \perp) : $x * y = x \prec y + x \succ y + x \perp y$ (post-algebra)

in a coherent way

[C.Bai, O. Bellier, L. Guo, X. Ni, 2011] \longrightarrow name of the procedure

[V. Gubarev, P.K., 2011]

The structures obtained are “dual” to di- and tri-algebras

Splitting an operad

Koszul duality for operads [V. Ginzburg, M. Kapranov, 1994]

\mathcal{P} binary quadratic operad

$$f \in \Omega \Rightarrow \nu(f) = 2, \quad \mathcal{P}(2) \curvearrowright S_2 \text{ space of binary operations}$$
$$\dim \mathcal{P}(2) < \infty$$

Defining identities have degree 3

$$\mathcal{P}(3) = \Omega(3)/V$$

$\Omega(3)$ = all polylinear terms of degree 3

$$S_3 \curvearrowright V \subset \Omega(3)$$

$$\Omega^\vee(3) = \Omega(3)^* \curvearrowright S_3$$

skew-transpose action:

$$\langle \varphi^\sigma, x \rangle = (-1)^\sigma \langle \varphi, x^\sigma \rangle$$

$V^\vee \subset \Omega^\vee(3)$ dual subspace

$$\mathcal{P}!(3) = \Omega^\vee(3)/V^\vee$$

$\mathcal{P}!$ Koszul-dual operad

Splitting an operad

for binary quadratic operads

$\text{Perm}^! = \text{pre-Lie}$

$\text{ComTrias}^! = \text{post-Lie}$ [B. Vallette, 2007]

\mathcal{P} binary quadratic operad

$$(\text{di-}\mathcal{P})^! = (\mathcal{P} \circ \text{Perm})^! = \mathcal{P}^! \bullet \text{Perm}^! = \mathcal{P}^! \bullet \text{pre-Lie}$$

$$(\text{tri-}\mathcal{P})^! = (\mathcal{P} \circ \text{ComTrias})^! = \mathcal{P}^! \bullet \text{ComTrias}^! = \mathcal{P}^! \bullet \text{post-Lie}$$

$\left. \begin{array}{l} \bullet \\ \circ \end{array} \right\}$ Black and white Manin products of binary quadratic operads

Splitting an operad

for binary quadratic operads

Com • pre-Lie = Zinb (Zinbiel algebras)

[J.-L. Loday, 1995]

As • pre-Lie = Dend (Dendriform algebras)

[J.-L. Loday, 2001]

Pois • pre-Lie = prePois (pre-Poisson algebras)

[M. Aguiar, 2000]

Dend • pre-Lie = Quad (Quadri-algebras)

[M. Aguiar, J.-L. Loday, 2004]

pre-Lie • pre-Lie = L-Dend (L-Dendriform algebras)

[C. Bai, L. Liu, X. Ni, 2010]

As • post-Lie = Tridend (Tridendriform algebras)

[J.-L. Loday, M. Ronco, 2004]

Com • post-Lie = CTD (Commutative tridendriform algebras)

[J.-L. Loday, 2007]

Splitting an operad

(Pre-algebras, general case)

$$\Omega = \{f_1, f_2, \dots\}$$

$$\nu(f_k) = n_k \geq 1$$

\mathcal{P} -Algebra:

$$(A, \Omega^A)$$

$$f_k^A : \underbrace{A \otimes \dots \otimes A}_{n_k} \rightarrow A$$

Defining identities Σ (multi-linear)

$$\Omega_2 = \{f_1^i, f_2^i, \dots\}$$

$$\nu(f_k^i) = n_k \geq 1$$

$$1 \leq i \leq n_k$$

pre- \mathcal{P} -algebra

$$(D, \Omega_2^D)$$

Defining identities **pre- Σ_2** :

$$P = \text{Perm}\langle x_1, x_2, \dots \rangle$$

$P \otimes D$ as an Ω -algebra:

$$f(p_1 \otimes a_1, \dots, p_n \otimes a_n) = \sum_{i=1}^n e_i^{(n)}(p_1, \dots, p_n) \otimes f^i(a_1, \dots, a_n)$$

Definition:

D is a pre- \mathcal{P} -algebra iff $P \otimes D$ is a \mathcal{P} -algebra.

$$D \models \text{pre-}\Sigma_2 \iff P \otimes D \models \Sigma$$

Splitting an operad

(Post-algebras, general case)

$$\Omega = \{f_1, f_2, \dots\}$$

$$\nu(f_k) = n_k \geq 1$$

\mathcal{P} -Algebra:

$$(A, \Omega^A)$$

$$f_k^A : \underbrace{A \otimes \dots \otimes A}_{n_k} \rightarrow A$$

Defining identities Σ (multi-linear)

$$\Omega_3 = \{f_1^H, f_2^H, \dots\}$$

$$\nu(f_k^H) = n_k \geq 1$$

$$H \subseteq \{1, \dots, n_k\}, H \neq \emptyset$$

post- \mathcal{P} -algebra

$$(D, \Omega_3^D)$$

Defining identities **post- Σ_3**

$$C = \text{ComTrias}\langle x_1, x_2, \dots \rangle$$

$C \otimes D$ as an Ω -algebra:

$$f(c_1 \otimes a_1, \dots, c_n \otimes a_n) = \sum_H e_H^{(n)}(c_1, \dots, c_n) \otimes f^H(a_1, \dots, a_n)$$

Definition:

D is a post- \mathcal{P} -algebra iff $C \otimes D$ is a \mathcal{P} -algebra.

$$D \models \text{pre-}\Sigma_3 \iff C \otimes D \models \Sigma$$

Splitting an operad

Theorem [V. Gubarev, P.K., 2011]

If \mathcal{P} is a binary quadratic operad then

$$(\text{pre-}\mathcal{P})^! = \text{di-}\mathcal{P}^! \quad (\text{post-}\mathcal{P})^! = \text{tri-}\mathcal{P}^!$$

Corollary

$$\text{pre-}\mathcal{P} = \mathcal{P} \bullet \text{pre-Lie}$$

$$\text{post-}\mathcal{P} = \mathcal{P} \bullet \text{post-Lie}$$

Splitting an operad

Corollary

If D is a (pre-)post- \mathcal{P} -algebra then $(C = \mathbb{k})$
for every (Perm-) ComTrias-algebra C
 $C \otimes D$ is a \mathcal{P} -algebra.

Example:

(1) $\mathcal{P} = \text{As}$, $\mu(x, y) = xy$

post-As = Tridend

$$\mu^1(x, y) = x \prec y, \mu^2(x, y) = x \succ y, \mu^{1,2}(x, y) = x \perp y$$

Then

$x * y = x \prec y + x \succ y + x \perp y$ is an associative product

(2) $\mathcal{P} = \text{Lie}$, $\mu(x, y) = [x, y]$

pre-Lie $\mu^1(x, y) = [x \prec y], \mu^2(x, y) = [x \succ y]$

$$[x \prec y] = -[y \succ x], \quad xy := [x \succ y] \quad \text{left-symmetric product}$$

Then

$$x * y = [x \succ y] + [x \prec y] = xy - yx \quad \text{is a Lie product}$$

Splitting an operad (Rota-Baxter operators)

Demo case:
one binary operation

$$\mu(x, y) = xy$$

Definition:

$R : A \rightarrow A$ *Rota-Baxter operator* of weight 0
if $R(x)R(y) = R(R(x)y + xR(y))$
for $x, y \in A$

[G. Baxter, 1960]
[G.-C. Rota, 1969]

Theorem (c.f. [M. Aguiar, 2000])

If A is a \mathcal{P} -algebra with a RB-operator R
then A w.r.t.

$$\mu^1(x, y) = xR(y), \quad \mu^2(x, y) = R(x)y$$

is a pre- \mathcal{P} -algebra.

Splitting an operad
(Rota-Baxter operators)

Demo case:
one binary operation

$$\mu(x, y) = xy$$

Definition:

$R : A \rightarrow A$ Rota-Baxter operator of weight 1
if $R(x)R(y) = R(R(x)y + xR(y) + xy)$
for $x, y \in A$

Theorem (c.f. [K. Ebrahimi-Fard, 2002])

If A is a \mathcal{P} -algebra with a RB-operator R of weight 1
then A w.r.t.

$$\mu^1(x, y) = xR(y), \quad \mu^2(x, y) = R(x)y, \quad \mu^{1,2}(x, y) = xy$$

is a post- \mathcal{P} -algebra.

Splitting an operad

(Rota-Baxter operators)

\mathcal{P} arbitrary operad

Theorem ([V. Gubarev, P.K., 2013])

If D is a (pre-) post- \mathcal{P} -algebra

then there exists a \mathcal{P} -algebra \widehat{D}

equipped with a Rota-Baxter operator R of weight $(0) 1$

such that D embeds into \widehat{D}

Splitting an operad

(Rota-Baxter operators)

\mathcal{P} arbitrary operad

Theorem ([V. Gubarev, P.K., 2013])

If D is a (pre-) post- \mathcal{P} -algebra

then there exists a \mathcal{P} -algebra \widehat{D}

equipped with a Rota-Baxter operator R of weight $(0) 1$

such that D embeds into \widehat{D}

Corollary (c.f. [K. Ebrahimi-Fard, L. Guo, 2008])

The universal enveloping Rota-Baxter \mathcal{P} -algebra

of a free (pre-)post- \mathcal{P} -algebra is free.

Splitting an operad

(PBW-type problems and special identities)

Splitting an operad

(PBW-type problems and special identities)



Splitting an operad

(PBW-type problems and special identities)

$\omega : \mathcal{R} \rightarrow \mathcal{P}$ morphism of operads

$$f(x_1, \dots, x_n) \in \mathcal{R}(n)$$

$$\omega(f)(x_1, \dots, x_n) \in \mathcal{P}(n)$$

$$C = \text{ComTrias}\langle c_1, c_2, \dots \rangle \quad (\text{or Perm})$$

Then

$$\omega(f)(c_1 \otimes x_1, \dots, c_n \otimes x_n) = \sum_{\substack{H \subseteq \{1, \dots, n\} \\ H \neq \emptyset}} e_H^{(n)}(c_1, \dots, c_n) \otimes g^H(x_1, \dots, g_n)$$

$$\boxed{\text{post-}\omega(f^H) := g^H} \quad (\text{or pre-}\omega)$$

Proposition.

$$\text{pre-}\omega: \text{pre-}\mathcal{R} \rightarrow \text{pre-}\mathcal{P}$$

$$\text{post-}\omega: \text{post-}\mathcal{R} \rightarrow \text{post-}\mathcal{P}$$

Splitting an operad

(PBW-type problems and special identities)

$$\omega : \mathcal{R} \rightarrow \mathcal{P}$$

$$\text{pre-}\omega : \text{pre-}\mathcal{R} \rightarrow \text{pre-}\mathcal{P}$$

$$\text{post-}\omega : \text{post-}\mathcal{R} \rightarrow \text{post-}\mathcal{P}$$

Theorem.

$S\text{pre-}\mathcal{R}\text{-algebras} \subseteq \text{pre-}S\mathcal{R}\text{-algebras}$

$S\text{post-}\mathcal{R}\text{-algebras} \subseteq \text{post-}S\mathcal{R}\text{-algebras}$

i.e., splitting of special identities

leads to special identities for pre- or post-algebras

[M. Bremner, S. Madariaga, 2013]: pre-Jord \rightarrow pre-As

No special identities of degree < 8

6 special identities of degree 8

Splitting an operad (A counterexample)

Example: ($\text{char } \mathbb{k} \neq 2$)

$\mathcal{R} = \text{Perm}$ $\mathcal{P} = \text{Com} + \text{Derivation } \partial: \partial^2(x) = 0$

$\omega: \mathcal{R} \rightarrow \mathcal{P}$ $xy = \partial(x)y$

$S\mathcal{R} = N3: x(yz) = (xy)z = 0$

Splitting:

pre- $N3$ is defined by

$$\begin{aligned} (x \prec y) \prec z &= 0 \\ (x \succ y) \prec z &= 0 \\ (x \prec y + x \succ y) \succ z &= 0 \\ x \prec (y \prec z + y \succ z) &= 0 \\ x \succ (y \prec z) &= 0 \\ x \succ (y \succ z) &= 0 \end{aligned}$$

pre- $\mathcal{P} = \text{Perm-algebra with a derivation } \partial, \partial^2 = 0$

pre- ω :

$$x \succ y = \partial(x)y, \quad x \prec y = y\partial(x)$$

Then

$$x \succ y + y \prec x = \partial(x)y + x\partial(y) = \partial(xy)$$

so

$$(x \succ y + y \prec x) \succ z = 0$$

holds on all special pre- \mathcal{R} -algebras

Thank you!