

LARGE DEVIATIONS OF SUMS OF RANDOM VARIABLES OF TWO TYPES

A. A. Borovkov

Let $\xi_1, \xi_2, \dots; \tau_1, \tau_2, \dots$ be two sequences of independent random variables, with ξ_i and τ_i distributed respectively as ξ and τ and with

$$\mathbb{E}|\xi| < \infty, \quad \mathbb{E}|\tau| < \infty, \quad S_n = \sum_{i=1}^n \xi_i, \quad T_m = \sum_{i=1}^m \tau_i.$$

In this article we study the asymptotics of large deviation probabilities of the sums $T_m + S_n$ for the following three classes of distribution tails for τ and ξ : regular (heavy), semiexponential, and exponentially decreasing. The numbers m and n may be either fixed or unboundedly increasing. The cause for appearance of this article is the articles [1, 2] addressing a particular case of the problem under consideration.

Key words and phrases: large deviation, sum of random variables, summands of two types.

Borovkov Aleksandr Alekseevich
Sobolev Institute of Mathematics,
630090 Novosibirsk, Russia.
E-mail: borovkov@math.nsc.ru

Received
April 17, 2001

Translated into English:

Siberian Advances in Mathematics, V. 11, N 4, 1–24 (2001).

1. Asmussen S., Klueppelberg G., and Sigman K. (1999) Sampling of sub-exponential times with queueing applications, *Stochastic Process. Appl.*, v. 79, N 2, 265–286.

2. Foss S. and Korshunov D. (2000) Sampling at random time with a heavy-tailed distribution, *Markov Process. Related Fields*, v. 6, N 4, 543–568.