

# THE TITS ALTERNATIVE FOR NON-SPHERICAL PRIDE GROUPS

NATALIA KOPTEVA AND GERALD WILLIAMS

ABSTRACT. Pride groups, or “groups given by presentations in which each defining relator involves at most two types of generators”, include Coxeter groups, Artin groups, triangles of groups, and Vinberg’s groups defined by periodic paired relations. We show that every non-spherical Pride group that is not a triangle of groups satisfies the Tits alternative.

## 1. INTRODUCTION

Pride groups, or “groups given by presentations in which each defining relator involves at most two types of generators” [11], include Coxeter groups, Artin groups, triangles of groups, and Vinberg’s groups defined by periodic paired relations. The cohomology of Pride groups was considered in [11], geometric invariants were considered in [8], and a Freiheitssatz was proven in [3].

In this paper we consider the Tits alternative for the class of Pride groups. Recall that a class of groups  $\mathcal{C}$  satisfies the *Tits alternative* if each group in  $\mathcal{C}$  contains a non-abelian free subgroup or has a soluble subgroup of finite index. This property is named after Tits who established that it is satisfied by the class of linear groups [14]; in particular, it holds for Coxeter groups.

The Tits alternative has been considered, for example, for the classes of mapping class groups of compact surfaces [7, 9], outer automorphism groups of free groups of finite rank [1, 2], subgroups of Gromov hyperbolic groups [5], groups acting on CAT(0) cubical complexes [12], triangles of groups [6], and groups defined by periodic paired relations [15, 16].

In this paper we prove the following

**Theorem 1.** *Every non-spherical Pride group  $G$  based on a graph with at least 4 vertices contains a non-abelian free subgroup, unless it is based on the graph shown in Figure 1, in which case  $G$  is virtually abelian and has presentation*

$$\langle x_1, x_2, x_3, x_4 \mid x_1^2, x_2^2, x_3^2, x_4^2, (x_1x_2)^2, (x_2x_3)^2, (x_3x_4)^2, (x_4x_1)^2 \rangle.$$

It is interesting to note that the “negatively curved” property of containing a non-abelian free subgroup is found in this non-positively curved class of groups.

We now give our formal definitions. Let  $\mathcal{G}$  be a finite simplicial graph with vertex set  $I(\mathcal{G})$ , and edge set  $E(\mathcal{G})$ . Further, let there be non-trivial groups  $G_i$  (with fixed finite presentations) associated to each vertex  $i \in I(\mathcal{G})$  and, in addition, for each edge  $\{i, j\} \in E(\mathcal{G})$  let  $R_{ij}$  be a finite collection of cyclically reduced words. We

---

*Date:* July 24, 2006.

*1991 Mathematics Subject Classification.* Primary: 20E05; Secondary: 20E07, 20F05, 20F06.

*Key words and phrases.* Pride group; Tits alternative; non-positively curved complex of groups.

The first author was partially supported by FP6 Marie Curie IIF Fellowship.




FIGURE 1

assume each word in  $R_{ij}$  is of free product length greater than or equal to 2 in  $G_i * G_j$ . The *Pride group* based on the graph  $\mathcal{G}$  with groups  $G_i$  assigned to the vertices and with edge relations  $R = \cup_{\{i,j\} \in E(\mathcal{G})} R_{ij}$  is the group  $G = *_{i \in I(\mathcal{G})} G_i / N$ , where  $N$  is the normal closure of  $R$  in  $*_{i \in I(\mathcal{G})} G_i$ .

We refer to the groups  $G_i$  as *vertex groups*, and we define the *edge groups* to be  $G_{ij} = \{G_i * G_j\} / N_{ij}$ , where  $\{i, j\} \in E(\mathcal{G})$  and where  $N_{ij}$  is the normal closure of  $R_{ij}$  in  $G_i * G_j$ . More generally, if  $\mathcal{F}$  is any full subgraph of  $\mathcal{G}$  with vertex set  $I(\mathcal{F}) \subseteq I(\mathcal{G})$ , then the *subgraph group*  $G_{\mathcal{F}}$  is  $\{*_{i \in I(\mathcal{F})} G_i\} / \{N_{ij} \mid \{i, j\} \in E(\mathcal{F})\}$ . In particular,  $G_{\mathcal{G}} = G$ .

For each  $i, j \in I(\mathcal{G})$ , the natural homomorphisms  $G_i \rightarrow G_{ij}$ ,  $G_j \rightarrow G_{ij}$  determine a homomorphism  $G_i * G_j \rightarrow G_{ij}$ . Let  $m_{ij}$  denote the length of a shortest non-trivial element in its kernel (in the usual length function on the free product), or put  $m_{ij} = \infty$  if the kernel is trivial. Note that either  $m_{ij} = 1$  (in which case one of the natural maps  $G_i \rightarrow G_{ij}$ ,  $G_j \rightarrow G_{ij}$  is not injective), or  $m_{ij}$  is even or infinite. The *Gersten-Stallings angle*  $(G_{ij}; G_i, G_j)$  between the groups  $G_i$  and  $G_j$  in the group  $G_{ij}$  is defined to be  $2\pi/m_{ij}$  for  $m_{ij} > 1$ , and 0 for  $m_{ij} = \infty$  [13].

In [11] Pride formulated the following asphericity condition. A Pride group is said to be *non-spherical* if

- (i)  $(G_{ij}; G_i, G_j) \leq \pi/2$  for all  $i, j \in I(\mathcal{G})$ ; and
- (ii) for any triangle  $\{i, j, k\}$  in  $\mathcal{G}$

$$(G_{ij}; G_i, G_j) + (G_{jk}; G_j, G_k) + (G_{ik}; G_i, G_k) \leq \pi.$$

In the non-spherical case we can assume that the graph  $\mathcal{G}$  is complete. To see this, observe that if  $i, j \in I(\mathcal{G})$  and  $\{i, j\} \notin E(\mathcal{G})$  then we can add the edge  $\{i, j\}$  and set  $R_{ij} = \emptyset$  without changing the group  $G$ .

If  $|I(\mathcal{G})| = 3$  then the Pride group  $G$  is the colimit of a triangle of groups. We also remark that every Pride group in which  $m_{ij} > 1$  for all  $i, j$  can be represented in terms of a 2-complex of groups. Moreover, if the Pride group is non-spherical then the corresponding complex can be chosen to be non-spherical.

## 2. PROOF OF THEOREM 1

Our method of prove has evolved from that developed in [4] and [6].

Let  $G = G_{\mathcal{G}}$  be a non-spherical Pride group, where  $\mathcal{G}$  is complete. First suppose that  $\mathcal{G}$  has four vertices. Let  $I(\mathcal{G}) = \{1, 2, 3, 4\}$  and let  $X = G_1$ ,  $Y = G_2$ ,  $Z = G_3$  and  $T = G_4$ . We shall sometimes write  $G_{XY}$  for  $G_{12}$ ,  $G_{XZ}$  for  $G_{13}$  and so on. Label the vertices of  $\mathcal{G}$  by the vertex groups and each edge  $\{i, j\}$  by  $(G_{ij}; G_i, G_j)$ .

If  $(G_{ij}; G_i, G_j) + (G_{jk}; G_j, G_k) + (G_{ik}; G_i, G_k) < \pi$  for some  $\{i, j, k\} \subset I(\mathcal{G})$  then, by [6],  $G_{ijk}$  contains a non-abelian free subgroup. By [3], every subgraph group embeds, so  $G$  also contains a non-abelian free subgroup. Hence, we may assume that for all  $i, j, k \in I(\mathcal{G})$  the angle sum is exactly  $\pi$ . Therefore, the edges that do not share any vertices have the same labels and all triangles in  $\mathcal{G}$  are labelled by one of  $\{\theta, \alpha, \beta\} = \{\pi/2, \pi/2, 0\}, \{\pi/2, \pi/3, \pi/6\}, \{\pi/2, \pi/4, \pi/4\}, \{\pi/3, \pi/3, \pi/3\}$ . Without loss of generality we may assume that  $\theta \geq \alpha \geq \beta$  and that

$$\begin{aligned} (G_{XZ}; G_X, G_Z) &= (G_{YT}; G_Y, G_T) = \theta, \\ (G_{XY}; G_X, G_Y) &= (G_{ZT}; G_Z, G_T) = \alpha, \\ (G_{YZ}; G_Y, G_Z) &= (G_{XT}; G_X, G_T) = \beta. \end{aligned}$$

Suppose  $(\theta, \alpha, \beta) \neq (\pi/2, \pi/2, 0)$  and consider a presentation  $\mathcal{P}$  for  $G$ . Since all the vertex groups are non-trivial, we may choose non-trivial elements  $x \in X$ ,  $y \in Y$ ,  $z \in Z$  and  $t \in T$  such that  $x, y, z, t$  are all generators of  $\mathcal{P}$ . We shall show that  $u = xyztxyz$  has infinite order in  $G$  and that  $t$  and  $u$  generate a free product.

Let  $w(t, u) = t^{p_1} u^{q_1} \dots t^{p_m} u^{q_m}$  or  $w(t, u) = u^{q_1}$ , where  $m \geq 1$  and each  $p_i, q_i \neq 0$ , and assume that  $w(t, u) = 1$  in  $G$ . Consider a van Kampen diagram  $K$  over  $\mathcal{P}$  whose boundary label is  $w(t, u)$ . Let  $D$  be an extremal disk of  $K$ . We divide  $D$  into  $G_{ij}$ -regions. If two  $G_{ij}$ -regions intersect at least at one edge, then we can amalgamate them into a single region. We continue in this way as often as possible, and so get a division of  $D$  into *maximal  $G_{ij}$ -regions*. (Note that the resulting division of  $D$  is not necessarily unique.)

By [3], the edge groups embed, so it can be assumed that the maximal regions are simply connected. Let  $\hat{D}$  be the resulting diagram. On the boundary of  $\hat{D}$  an edge of  $\hat{D}$  is defined to be a longest path whose edges are labelled by elements of the same vertex group. In the interior an edge is defined to be the intersection of two adjacent maximal  $G_{ij}$ - and  $G_{ik}$ -regions. Note that it is a path labelled by elements of  $G_i$ .

Now place  $\hat{D}$  on the sphere and take its dual  $D^*$ . Let  $v_0$  be the vertex corresponding to  $\mathbb{S}^2 \setminus \hat{D}$ . We call a region of  $D^*$  *exterior* if it involves  $v_0$  and *interior* otherwise. We give each corner at a vertex of  $D^*$  of degree  $\delta$  the angle  $2\pi/\delta$ . The curvature  $c(\Delta)$  of a region  $\Delta$  of degree  $q$  whose vertices have degrees  $\delta_1, \delta_2, \dots, \delta_q$  is then defined by

$$c(\Delta) = (2 - q)\pi + \sum_{i=1}^q \frac{2\pi}{\delta_i}.$$

Then

$$\sum_{\Delta \subset D^*} c(\Delta) = 2\pi\chi(\mathbb{S}^2) = 4\pi.$$

Let  $\Delta$  be an interior region of  $D^*$  of degree  $q$ . Observe that  $q \geq 3$ . If  $q \geq 4$  then

$$c(\Delta) \leq (2 - q)\pi + \sum_{i=1}^q \frac{2\pi}{4} \leq 0$$

and if  $q = 3$  then (for some distinct  $i, j, k \in I(\mathcal{G})$ )

$$c(\Delta) \leq -\pi + (G_{ij}; G_i, G_j) + (G_{jk}; G_j, G_k) + (G_{ik}; G_i, G_k) = 0.$$

Thus, the sum of the curvatures of interior regions is non-positive.

Consider exterior regions. Observe that an exterior region can be a 2-gon. It is convenient to define  $d(\Delta) = c(\Delta) - 2\pi/N$ , where  $N$  is the number of exterior regions. Then

$$\sum_{\text{ext. } \Delta} c(\Delta) = \sum_{\text{ext. } \Delta} d(\Delta) + 2\pi.$$

We aim to show  $\sum_{\text{ext. } \Delta} d(\Delta) \leq 0$  and obtain a contradiction.

Split the boundary of  $\widehat{D}$  into  $t^{p_i}$  and  $u^\varepsilon = (xyztxyz)^\varepsilon$  pieces, where  $\varepsilon = \pm 1$ . We now consider the sum of the curvatures of the exterior regions of  $D^*$  arising from each  $u^\varepsilon$  piece.




FIGURE 2

The analysis of a  $u^\varepsilon$  piece is essentially the same for both  $\varepsilon = +1$  and  $\varepsilon = -1$ , so we introduce the following notation. Let  $(a, b) = (x, z)$ ,  $(A, B) = (X, Z)$ ,  $(\phi, \psi) = (\alpha, \beta)$  if  $\varepsilon = +1$  and let  $(a, b) = (z^{-1}, x^{-1})$ ,  $(A, B) = (Z, X)$ ,  $(\phi, \psi) = (\beta, \alpha)$  if  $\varepsilon = -1$ . Note that with this convention  $\theta \geq \phi$  and  $\theta \geq \psi$ . Figure 2(a) indicates the Gersten-Stallings angles between the vertex groups and Figure 2(b) shows the form of a  $u^\varepsilon$  piece, where the exterior regions of  $D^*$  are labelled by  $\Delta_i$  ( $1 \leq i \leq 7$ ) and each  $\Delta_i$  is a  $q_i$ -gon.

Observe that at most four of the  $\Delta_i$  can be 2-gons. Denote the chain  $\Delta_1 \Delta_2 \dots \Delta_7$  by  $S$  and write  $d(S) = \sum_{i=1}^7 d(\Delta_i)$ . Denote by  $v_1$  the vertex of  $\Delta_1 \setminus \Delta_2$  adjacent to  $v_0$ .

We shall make frequent use of the following trivial observations. Let  $\Delta$  be an exterior  $q$ -gon. If  $q = 2$  then  $d(\Delta) \leq \pi/2$ . If  $q = 3$  then no two adjacent vertices of  $\Delta$  arise from maximal  $G_{AB}$ - or  $G_{YT}$ -regions, and so  $d(\Delta) \leq -\pi + \pi/2 + \pi/3 = -\pi/6$ . Similarly, if  $q \geq 4$  then  $d(\Delta) \leq -2\pi/3$ .

**Claim 1.** *If  $v_1$  does not arise from any maximal  $G_{AT}$ -region then  $d(S) \leq 0$ .*

*Proof.* Note that if  $d(\Delta_i) > \pi/3$  then  $i = 7$ . Hence, if  $|\{i \mid q_i = 2\}| \leq 2$  then  $d(S) \leq \pi/2 + \pi/3 - 5\pi/6 = 0$ .

Suppose that  $|\{i \mid q_i = 2\}| = 3$ . If  $q_i \geq 4$  for some  $i$  then  $d(S) \leq 2\pi/3 + \pi/2 + 3(-\pi/6) + (-2\pi/3) = 0$ . Hence, we may assume that  $q_i \leq 3$  for  $1 \leq i \leq 7$ . However, if  $q_i = q_{i+2} = 2$  for any  $i \leq 4$  then  $q_{i+1} \geq 4$ ; moreover, if  $q_7 \neq 2$  then this condition holds for some  $i$ . Thus  $\{i \mid q_i = 2\} = \{1, 4, 7\}$ ,  $\{1, 5, 7\}$ ,  $\{2, 5, 7\}$ . Label consecutive vertices of  $S$  by  $v_2, \dots, v_5$  so that  $v_2$  is adjacent to  $v_1$ .

**{1, 4, 7}.** The vertices  $v_1$  and  $v_3$  arise from maximal  $G_{AY}$ - and  $G_{AT}$ -regions, respectively. Moreover, at most two of the five angles in  $\Delta_5$ ,  $\Delta_6$ , and  $\Delta_7$  can be greater than  $\pi/3$ . Then

$$d(S) \leq \phi + (\phi + \theta - \pi) + (\theta + \psi - \pi) + \psi + 3\frac{\pi}{3} + 2\frac{\pi}{2} - 2\pi = 2\phi + 2\psi + 2\theta - 2\pi = 0.$$

**{1, 5, 7}.** Since  $v_1$  and  $v_4$  arise from maximal  $G_{AY}$ -regions and  $q_2 = q_3 = q_4 = 3$ , either  $v_2$  arises from a maximal  $G_{AB}$ -region and  $v_3$  arises from a maximal  $G_{AT}$ -region, or  $v_2$  arises from a maximal  $G_{YB}$ -region and  $v_3$  arises from a maximal  $G_{YT}$ -region, see Figure 3(a). In both cases

$$d(S) \leq 5\phi + 2\psi + 4\theta - 4\pi = 3\phi + 2\theta - 2\pi \leq 0.$$

**{2, 5, 7}.** Since  $v_2$  and  $v_4$  arise from maximal  $G_{BY}$ - and  $G_{AY}$ -regions, we immediately get

$$d(S) \leq 3\phi + 3\psi + 5\theta - 4\pi = 2\theta - \pi \leq 0.$$

Finally, suppose that  $|\{i \mid q_i = 2\}| = 4$ . Then  $\{i \mid q_i = 2\} = \{1, 3, 5, 7\}$  and hence  $q_2 \geq 4$  and  $q_4 \geq 4$ . Since  $d(\Delta_i) \leq \phi$  for  $i = 1, 3, 5$  and  $d(\Delta_7) \leq \pi/2$ , we get  $d(S) \leq 3\phi + \pi/2 - 4\pi/3 - \pi/6 \leq 0$ .  $\square$




FIGURE 3

**Claim 2.** If  $v_1$  arises from a maximal  $G_{AT}$ -region then  $d(S) \leq -\pi/3$ .

*Proof.* Since  $v_1$  arises from a maximal  $G_{AT}$ -region,  $\Delta_1$  is not a 2-gon and, therefore,  $|\{i \mid q_i = 2\}| \leq 3$ . If  $|\{i \mid q_i = 2\}| \leq 1$  then  $d(S) \leq \pi/2 + 6(-\pi/6) = -\pi/2$ .

Suppose that  $|\{i \mid q_i = 2\}| = 2$ . If  $q_i \geq 4$  for some  $i$  then  $d(S) \leq -\pi/2$ . Hence, we may assume that  $q_i \leq 3$  for  $1 \leq i \leq 7$ . However, if  $q_i = q_{i+2} = 2$  for any  $i \leq 4$  then  $q_{i+1} \geq 4$ ; moreover, if  $q_2 = 2$  then  $q_1 \geq 4$ . This reduces us to the four cases:  $\{i \mid q_i = 2\} = \{3, 6\}$ ,  $\{3, 7\}$ ,  $\{4, 7\}$ ,  $\{5, 7\}$ . Label consecutive vertices of  $S$  by  $v_2, \dots, v_6$  so that  $v_2$  is adjacent to  $v_1$ .

**{3, 6}.** Since  $v_3$  arises from a maximal  $G_{BT}$ -region and  $v_5$  arises from a maximal  $G_{YB}$ -region, we have  $d(S) \leq 4\psi + 3\phi + 5\theta - 5\pi \leq -2\pi/3$ .

**{3, 7}.** If  $u^\epsilon$  is followed by  $t^{p_i}$ , then both  $v_3$  and  $v_6$  arise from maximal  $G_{BT}$ -regions and thus

$$\begin{aligned} d(S) &= d(\Delta_1\Delta_2\Delta_3) + d(\Delta_4\Delta_5\Delta_6) + d(\Delta_7) \\ &\leq (\psi + 2\phi + 2\theta - 2\pi) - 3\frac{\pi}{6} + \phi \leq -\frac{\pi}{3}. \end{aligned}$$

So suppose that  $u^\epsilon$  is followed by  $a$  (see Figure 3(b)). Then  $v_6$  arises from a maximal  $G_{AB}$ -region. If  $v_4$  arises from a maximal  $G_{AT}$ -region or  $v_5$  arises from a maximal  $G_{BY}$ -region then  $d(S) \leq 3\psi + 3\phi + 6\theta - 5\pi \leq -\pi/2$ . Hence, we may assume that

$v_4$  arises from a maximal  $G_{AB}$ -region and  $v_5$  arises from a maximal  $G_{AY}$ -region. It follows that  $\deg(v_4) \geq 5$ , that is, the angle at  $v_4$  is at most  $2\pi/5$ , so we have

$$\begin{aligned} d(S) &= d(\Delta_1\Delta_2\Delta_3) + d(\Delta_4\Delta_5) + d(\Delta_6\Delta_7) \\ &\leq (\psi + 2\phi + 2\theta - 2\pi) + \left(2\phi + 2\frac{2\pi}{5} - 2\pi\right) + \left(-\frac{\pi}{6} + \frac{\pi}{2}\right) \leq -\frac{11\pi}{30}. \end{aligned}$$

**{4, 7}.** Since both  $v_1$  and  $v_4$  arise from maximal  $G_{AT}$ -regions,  $d(\Delta_1\Delta_2\Delta_3) \leq 2\psi + 2\phi + 2\theta - 3\pi = -\pi$ . Then  $d(S) = d(\Delta_1\Delta_2\Delta_3) + d(\Delta_4\Delta_5\Delta_6\Delta_7) \leq -\pi + \psi - 2\pi/6 + \pi/2 \leq -\pi/2$ .

**{5, 7}.** If  $u^\varepsilon$  is followed by  $t^{p_i}$ , then  $q_5 \geq 4$ , a contradiction. Suppose  $u^\varepsilon$  is followed by  $a$ . The vertex  $v_5$  arises from a maximal  $G_{AY}$ -region and  $v_6$  arises from a maximal  $G_{AB}$ -region. If  $v_3$  arises from a maximal  $G_{BY}$ -region or  $v_4$  arises from a maximal  $G_{AT}$ -region then  $d(S) \leq 3\psi + 3\phi + 6\theta - 5\pi \leq -\pi/2$ . Therefore, we may assume that  $v_4$  arises from a maximal  $G_{YT}$ -region. It follows that  $v_3$  arises from a maximal  $G_{BT}$ -region and hence  $\deg(v_4) \geq 5$ . Thus,

$$d(S) \leq \psi + 5\phi + 4\theta + 2\frac{2\pi}{5} - 5\pi \leq -\frac{11\pi}{30}.$$

Now suppose that  $|\{i \mid q_i = 2\}| = 3$ . Then  $\{i \mid q_i = 2\}$  is one of the following:  $\{2, 4, 6\}$ ,  $\{2, 4, 7\}$ ,  $\{2, 5, 7\}$ ,  $\{3, 5, 7\}$ . However, if  $q_2 = q_4 = 2$  then  $q_1 \geq 4$  and  $q_3 \geq 4$ , and so  $d(S) \leq -\pi/2$ .

**{2, 5, 7}.** Since  $q_1 \geq 4$ , we may assume that  $q_i \leq 3$  for all  $1 < i \leq 7$ . In particular,  $q_6 = 3$ , so the 2-gon  $\Delta_7$  comes from a maximal  $G_{AB}$ -region. Since  $\Delta_2$  and  $\Delta_5$  come from maximal  $G_{BY}$ - and  $G_{AY}$ -regions, respectively, we have

$$d(S) \leq 3\phi + 4\psi + 5\theta - 5\pi = \psi + 2\theta - 2\pi \leq -\frac{2\pi}{3}.$$

**{3, 5, 7}.** Then  $q_4 \geq 4$  and again we may assume that  $q_i \leq 3$  for all  $i \neq 4$  and therefore the 2-gon  $\Delta_7$  comes from a maximal  $G_{AB}$ -region. Since  $\Delta_3$  and  $\Delta_5$  come from maximal  $G_{BT}$ - and  $G_{AY}$ -regions, respectively, we have

$$d(S) \leq 6\phi + \psi + 5\theta - 5\pi = 5\phi + 4\theta - 4\pi \leq -\frac{\pi}{3}.$$

□

If  $w(t, u) = u^{q_1}$  then Claims 1 and 2 imply the required contradiction that  $\sum_{ext. \Delta} d(\Delta) \leq 0$ . Hence,  $u$  has infinite order in  $G$ .

Now suppose  $w(t, u) = t^{p_1}u^{q_1} \dots t^{p_m}u^{q_m}$ . For each syllable  $t^{p_i}u^{q_i} = t^{p_i}u^\varepsilon u^{q_i - \varepsilon}$ , consider the part of the boundary corresponding to  $t^{p_i}u^\varepsilon$ . Label the first exterior region  $\Delta_0$  and, as before, label the remaining regions  $\Delta_1, \dots, \Delta_7$ .

If  $q_0 \neq 2$  then  $d(\Delta_0) \leq -\pi/6$  and, therefore,  $d(\Delta_0 S) = d(\Delta_0) + d(S) < 0$ . If  $\Delta_0$  is a 2-gon then  $v_1$  arises from a  $G_{AT}$ -region. But then, by Claim 2,  $d(S) \leq -\pi/3$  and, hence,  $d(\Delta_0 S) \leq \psi - \pi/3 \leq 0$ . It follows that the sum of the  $d$ -values of all exterior regions arising from any syllable  $t^{p_i}u^{q_i}$  is non-positive, and the required contradiction follows. Hence,  $t$  and  $u$  generate a free product in  $G$  and, since  $u$  has infinite order,  $G$  contains a non-abelian free subgroup.

Now suppose  $(\theta, \alpha, \beta) = (\pi/2, \pi/2, 0)$ . Then  $G$  is isomorphic to an amalgamated free product  $L *_K M$ , where  $L = G_{14} *_K G_{34}$ ,  $M = G_{12} *_K G_{23}$  and  $K = G_1 *_K G_3$ . We may assume that  $|G_1| = |G_3| = 2$  for otherwise  $K$  (and hence  $G$ ) contains

a non-abelian free subgroup. Similarly we may assume that  $|G_2| = |G_4| = 2$ , so  $G_i = \langle x_i | x_i^2 \rangle$  for all  $i$ . Further, since each  $m_{i,i+1} = 4$ ,  $\{(x_i x_{i+1})^2\} \subseteq R_{i,i+1}$ . Hence  $G_{i,i+1} = \langle x_i, x_{i+1} | x_i^2, x_{i+1}^2, (x_i x_{i+1})^2 \rangle \cong D_4$ . Therefore,  $G$  has presentation

$$\langle x_1, x_2, x_3, x_4 | x_1^2, x_2^2, x_3^2, x_4^2, (x_1 x_2)^2, (x_2 x_3)^2, (x_3 x_4)^2, (x_4 x_1)^2 \rangle.$$

Since  $G$  is a group of isometries of the Euclidean plane, it is virtually abelian.

Thus the theorem is proved when  $G$  is based on a graph with four vertices. To complete the proof in the general case, it remains to note that when  $\mathcal{G}$  has five or more vertices some subgraph of  $\mathcal{G}$  with four vertices necessarily gives rise to a group that contains a non-abelian free subgroup. Since by [3] subgraph groups embed, Theorem 1 is proved.

### 3. APPLICATION

We consider the following class of groups which generalizes the groups defined by periodic paired relations [15]. Let  $n \geq 3$ ,  $1 \leq i, j \leq n$ ,  $n_{ij} \geq 1$  and  $1 \leq t \leq n_{ij}$ . For each such  $i, j, t$  let  $2 \leq q_i, q_{i,j;t} \leq \infty$  and suppose  $w_{i,j;t}(x_i, x_j)$  is a cyclically reduced word in  $x_i$  and  $x_j$ . Define

$$\Gamma = \langle x_1, \dots, x_n | x_i^{q_i}, w_{i,j;t}(x_i, x_j)^{q_{i,j;t}} (1 \leq i, j \leq n, 1 \leq t \leq n_{ij}) \rangle.$$

Each group  $\Gamma$  can be realized as a Pride group by setting  $G_i = \langle x_i | x_i^{q_i} \rangle$  and  $R_{ij} = \{w_{i,j;t} | 1 \leq t \leq n_{ij}\}$ . For each  $i, j$  define  $r_{ij} = \min\{\ell_{i,j;t} q_{i,j;t} | 1 \leq t \leq n_{ij}\}$ , where  $\ell_{i,j;t}$  denotes the free product length of  $w_{i,j;t}$ . If  $1/r_{ij} + 1/r_{jk} + 1/r_{ik} \leq 1/2$  for all distinct  $1 \leq i, j, k \leq n$  then by the Spelling Theorem for generalized triangle groups [6], the Pride group  $\Gamma$  is non-spherical.

**Corollary.** *Let  $\Gamma$  be as defined above with  $n \geq 4$ . If  $1/r_{ij} + 1/r_{jk} + 1/r_{ik} \leq 1/2$  for all distinct  $i, j, k$ , then  $\Gamma$  contains a non-abelian free subgroup unless it has presentation*

$$\langle x_1, x_2, x_3, x_4 | x_1^2, x_2^2, x_3^2, x_4^2, (x_1 x_2)^2, (x_2 x_3)^2, (x_3 x_4)^2, (x_4 x_1)^2 \rangle,$$

*in which case  $\Gamma$  is virtually abelian.*

*Acknowledgements.* The second author would like to thank the department of mathematics at the Université de Provence for its hospitality during a research visit in July 2006, when part of this work was carried out.

### REFERENCES

- [1] M. Bestvina, M. Feighn and M. Handel, *The Tits alternative for  $\text{Out}(F_n)$ . I. Dynamics of exponentially-growing automorphisms*, Ann. of Math. (2) **151** (2000), no. 2, 517–623.
- [2] M. Bestvina, M. Feighn and M. Handel, *The Tits alternative for  $\text{Out}(F_n)$ . II. A Kolchin type theorem*, Ann. of Math. (2) **161** (2005), no. 1, 1–59.
- [3] J. M. Corson, *Amalgamated sums of groups*, Proc. Edinburgh Math. Soc. (2) **39** (1996), no. 3, 561–570.
- [4] M. Edjvet, G. Rosenberger, M. Stille and R. M. Thomas, *On certain finite generalized tetrahedron groups*, Computational and geometric aspects of modern algebra (Edinburgh, 1998), 54–65, London Math. Soc. Lecture Note Ser., **275**, Cambridge Univ. Press, Cambridge, 2000.
- [5] M. Gromov, *Hyperbolic groups*, Essays in group theory, 75–263, Math. Sci. Res. Inst. Publ., 8, Springer, New York, 1987.
- [6] J. Howie and N. Kopteva, *The Tits alternative for generalized tetrahedron groups*, J. Group Theory, **9** (2006), no. 2, 173–189.
- [7] N. V. Ivanov, *Algebraic properties of the Teichmüller modular group*, (Russian) Dokl. Akad. Nauk SSSR **275** (1984), no. 4, 786–789.

- [8] J. Meier, *Geometric invariants for Artin groups*, Proc. London Math. Soc. (3) **74** (1997), no. 1, 151–173.
- [9] J. McCarthy, A “Tits-alternative” for subgroups of surface mapping class groups, Trans. Amer. Math. Soc. **291** (1985), no. 2, 583–612.
- [10] G. Noskov and E. B. Vinberg, *Strong Tits alternative for subgroups of Coxeter groups*, J. Lie Theory **12** (2002), no. 1, 259–264.
- [11] S. J. Pride, *The (co)homology of groups given by presentations in which each defining relator involves at most two types of generators*, J. Austral. Math. Soc. Ser. A **52** (1992), no. 2, 205–218.
- [12] M. Sageev and D. T. Wise, *The Tits alternative for CAT(0) cubical complexes*, Bull. London Math. Soc. **37** (2005), no. 5, 706–710.
- [13] J. R. Stallings, *Non-positively curved triangles of groups*, Group Theory from a Geometrical Viewpoint (Trieste, 1990, ed. E. Ghys, A. Haefliger and P. de la Harpe), World Scientific, Singapore (1991), pp. 491–503.
- [14] J. Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270.
- [15] E. B. Vinberg, *Groups defined by periodic paired relations*, Sb. Math. **188** (1997), no. 9, 1269–1278.
- [16] G. Williams, *The Tits alternative for groups defined by periodic paired relations*, Comm. Algebra **34** (2006), no. 1, 251–258.

LATP (UMR CNRS 6632), CMI, 39 RUE F. JOLIOT CURIE, 13453 MARSEILLE CEDEX 13, FRANCE

*E-mail address:* `kopteva@cmi.univ-mrs.fr`

INSTITUTE OF MATHEMATICS, STATISTICS, AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT, CANTERBURY, KENT CT2 7NF, UK

*E-mail address:* `g.williams@kent.ac.uk`